

1P-344-C100

Monoclonal Antibody to TRIM Phycoerythrin (PE) conjugated (0.1 mg)

Clone:	TRIM-04
Isotype:	Mouse IgG2a
Specificity:	The antibody TRIM-04 reacts with T cell receptor-interacting molecule (TRIM), a 30 kDa adaptor protein expressed by T cells.
Regulatory Status:	RUO
Immunogen:	Recombinant intracellular domain (aa 29-186) of human TRIM.
Species Reactivity:	Human
Negative Species:	Mouse
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis (see application note on www.exbio.cz). Suggested working dilution is 1:50. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	TRIM (T cell receptor-interacting molecule), also known as TRAT1 (T cell receptor associated transmembrane adaptor 1) is a 30 kDa protein expressed by T cells as a cystein-linked homodimer. It associates with TCR-CD3-zeta-chain complex and becomes phosphorylated by Src-family kinases. TRIM is potentially involved in negative regulation of TCR-mediated signaling, but its role remains unclear.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Huynh T, Wurch A, Bruyns E, Korinek V, Schraven B, Eichmann K.: Developmentally regulated expression of the transmembrane adaptor protein trim in fetal and adult T cells. *Scand J Immunol.* 2001 Jul-Aug;54(1-2):146-54.
 - *Horejsí V, Zhang W, Schraven B.: Transmembrane adaptor proteins: organizers of immunoreceptor signalling. *Nat Rev Immunol.* 2004 Aug;4(8):603-16.
 - *Simeoni L, Smida M, Posevitz V, Schraven B, Lindquist JA.: Right time, right place: the organization of membrane proximal signaling. *Semin Immunol.* 2005 Feb;17(1):35-49.
 - *Tedoldi S, Paterson JC, Hansmann ML, Natkunam Y, Rüdiger T, Angelisova P, Du MQ, Robertson H, Roncador G, Sanchez L, Pozzobon M, Masir N, Barry R, Pileri S, Mason DY, Marafioti T, Horejsí V.: Transmembrane adaptor molecules: a new category of lymphoid-cell markers. *Blood.* 2006 Jan 1;107(1):213-21.
 - *Kirchgessner H, Dietrich J, Scherer J, Isomaki P, Korinek V, Hilgert I, Bruyns E, Leo A, Cope AP, Schraven B.: The transmembrane adaptor protein TRIM regulates T cell receptor (TCR) expression and TCR-mediated signaling via an association with the TCR zeta chain. *J Exp Med.* 2001 Jun 4;193(11):1269-84.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz