



1P-413-T100

Monoclonal Antibody to CD10 Phycoerythrin (PE) conjugated (100 tests)

Clone:	LT10
Isotype:	Mouse IgG1
Specificity:	The antibody LT10 reacts with CD10 antigen (CALLA - Common acute lymphatic leukemia antigen), a 100 kDa type II integral membrane protein.
Regulatory Status:	RUO
Immunogen:	mouse NALM-6 leukemia pre-B cell line (tissue/cell preparation)
Species Reactivity:	Human, Other not tested
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD10 (neutral endopeptidase – NEP, common acute lymphocytic leukemia antigen – CALLA, membrane metallo-endopeptidase – MME, enkephalinase) is a 100-kDa cell surface zinc metalloprotease cleaving peptide bonds on the N-terminus of hydrophobic amino acids and inactivating multiple physiologically active peptides. CD10 is expressed on various normal cell types, including lymphoid precursor cells, germinal center B lymphocytes, and some epithelial cells, and its expression level serves as a marker for diagnostics of many carcinomas. CD10 is also a differentiation antigen for early B-lymphoid progenitors in the B-cell differentiation pathway and has a key role in regulation of growth, differentiation and signal transduction of many cellular systems.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References:

*Suzuki T, Ino K, Kikkawa F, Uehara C, Kajiyama H, Shibata K, Mizutani S: Neutral endopeptidase/CD10 expression during phorbol ester-induced differentiation of choriocarcinoma cells through the protein kinase C- and extracellular signal-regulated kinase-dependent signalling pathway. *Placenta*. 2002 Jul;23(6):475-82.

*Yada K, Kashima K, Daa T, Kitano S, Fujiwara S, Yokoyama S: Expression of CD10 in basal cell carcinoma. *Am J Dermatopathol*. 2004 Dec;26(6):463-71.

*Braham H, Trimeche M, Ziadi S, Mestiri S, Mokni M, Amara K, Hachana M, Sriha B, Korbi S: CD10 expression by fusiform stromal cells in nasopharyngeal carcinoma correlates with tumor progression. *Virchows Arch*. 2006 Aug;449(2):220-4.

*Dall'Era MA, True LD, Siegel AF, Porter MP, Sherertz TM, Liu AY: Differential expression of CD10 in prostate cancer and its clinical implication. *BMC Urol*. 2007 Mar 2;7:3.

*Leukocyte Typing VII., Mason D. et al. (Eds.), Oxford University Press (2002).

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz