



Antibodies

1P-492-T025

## Monoclonal Antibody to CD2 Phycoerythrin (PE) conjugated (25 tests)

<b>Clone:</b>	LT2
<b>Isotype:</b>	Mouse IgG2b
<b>Specificity:</b>	The antibody LT2 reacts with CD2, a 50 kDa glycoprotein present on the human peripheral blood T lymphocytes and NK cells; also expressed by all thymocytes. HLDA VI; WS Code T 6T-008
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Normal human blood lymphocytes.
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.5 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD2 belongs to T lymphocyte glycoproteins of immunoglobulin superfamily. Its interaction with CD58 stabilizes adhesion between T cells and antigen presenting or target cells. Relatively low affinity of CD2 to CD58 (as measured in solution) is compensated within the two-dimensional cell-cell interface to provide tight adhesion. Moreover, T cell activation induces increased CD2 expression and its lateral mobility, making easier contact between CD2 and CD58. Subsequently, T cell activation causes fixation of CD58-CD2 at sites of cell-cell contact, thereby strengthening intercellular adhesion. CD2 deficiency reduces intestinal inflammation and helps to control infection.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*Dustin ML: Adhesive bond dynamics in contacts between T lymphocytes and glass-supported planar bilayers reconstituted with the immunoglobulin-related adhesion molecule CD58. *J Biol Chem.* 1997 Jun 20;272(25):15782-8.
  - \*Zhu DM, Dustin ML, Cairo CW, Thatte HS, Golan DE.: Mechanisms of Cellular Avidity Regulation in CD2-CD58-Mediated T Cell Adhesion. *ACS Chem Biol.* 2006 Nov 21;1(10):649-58.
  - \*Pawlowski NN, Struck D, Grollich K, Kuhl AA, Zeitz M, Liesenfeld O, Hoffmann JC.: CD2 deficiency partially prevents small bowel inflammation and improves parasite control in murine *Toxoplasma gondii* infection. *World J Gastroenterol.* 2007 Aug 21;13(31):4207-13.
  - Filatov AV, Bachurin PS, Markova NA, Surnakova NE: [A panel of monoclonal antibodies against human lymphocyte antigens] *Eksp Onkol.* 1989;11(2):28-32.
  - \*Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
  - \*Gannon GA, Rhind SG, Suzui M, Zamecnik J, Sabiston BH, Shek PN, Shephard RJ: beta-Endorphin and natural killer cell cytolytic activity during prolonged exercise. Is there a connection? *Am J Physiol.* 1998 Dec;275(6 Pt 2):R1725-34.
  - \*Lin CW, Liu TY, Chen SU, Wang KT, Medeiros LJ, Hsu SM: CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. *Blood.* 2005 Nov 15;106(10):3567-74.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)