

1P-595-C100

Monoclonal Antibody to CD11b (mouse) Phycoerythrin (PE) conjugated (0.1 mg)

Clone:	M1/70
Isotype:	Rat IgG2b
Specificity:	The rat monoclonal antibody M1/70 detects CD11b (integrin alphaM subunit), a type I transmembrane protein mainly expressed on monocytes/macrophages, granulocytes and NK-cells, which associates with CD18 to form Mac-1 integrin that plays important role in cell-cell interactions.
Regulatory Status:	RUO
Immunogen:	B10 mouse spleen cells enriched for T cells
Species Reactivity:	Human, Non-Human Primates, Mouse, Rabbit
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working concentration is 0.5 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Springer T, Galfrè G, Secher DS, Milstein C: Monoclonal xenogeneic antibodies to murine cell surface antigens: identification of novel leukocyte differentiation antigens. *Eur J Immunol.* 1978 Aug;8(8):539-51.
- *Ault KA, Springer TA: Cross-reaction of a rat-anti-mouse phagocyte-specific monoclonal antibody (anti-Mac-1) with human monocytes and natural killer cells. *J Immunol.* 1981 Jan;126(1):359-64.
- *Sanchez-Madrid F, Simon P, Thompson S, Springer TA: Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. *J Exp Med.* 1983 Aug 1;158(2):586-602.
- *Whiteland JL, Nicholls SM, Shimeld C, Easty DL, Williams NA, Hill TJ: Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. *J Histochem Cytochem.* 1995 Mar;43(3):313-20.
- *Dembic Z, Schenck K, Bogen B: Dendritic cells purified from myeloma are primed with tumor-specific antigen (idiotype) and activate CD4+ T cells. *Proc Natl Acad Sci U S A.* 2000 Mar 14;97(6):2697-702.
- *Welt FG, Edelman ER, Simon DI, Rogers C: Neutrophil, not macrophage, infiltration precedes neointimal thickening in balloon-injured arteries. *Arterioscler Thromb Vasc Biol.* 2000 Dec;20(12):2553-8.
- *Zhang Y, McCormick LL, Desai SR, Wu C, Gilliam AC: Murine sclerodermatous graft-versus-host disease, a model for human scleroderma: cutaneous cytokines, chemokines, and immune cell activation. *J Immunol.* 2002 Mar 15;168(6):3088-98.
- *Brickson S, Ji LL, Schell K, Olabisi R, St Pierre Schneider B, Best TM: M1/70 attenuates blood-borne neutrophil oxidants, activation, and myofiber damage following stretch injury. *J Appl Physiol.* 2003 Sep;95(3):969-76.
- *Leon F, Contractor N, Fuss I, Marth T, Lahey E, Iwaki S, la Sala A, Hoffmann V, Strober W, Kelsall BL: Antibodies to complement receptor 3 treat established inflammation in murine models of colitis and a novel model of psoriasiform dermatitis. *J Immunol.* 2006 Nov 15;177(10):6974-82.
- *Takagi H, Numazaki M, Kajiwara T, Abe Y, Ishii M, Kato C, Kojima N: Cooperation of specific ICAM-3 grabbing nonintegrin-related 1 (SIGIRR) and complement receptor type 3 (CR3) in the uptake of oligomannose-coated liposomes by macrophages. *Glycobiology.* 2009 Mar;19(3):258-66.
- *And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz