

1P-596-C025

Monoclonal Antibody to CD19 (mouse) Phycoerythrin (PE) conjugated (0.025 mg)

Clone:	1D3
Isotype:	Rat IgG2a
Specificity:	The rat monoclonal antibody 1D3 detects mouse CD19, 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.
Regulatory Status:	RUO
Immunogen:	Mouse CD19-transfected cell line
Species Reactivity:	Mouse
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working dilution is 1 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell threshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References:

*Krop I, Shaffer AL, Fearon DT, Schlissel MS: The signaling activity of murine CD19 is regulated during cell development. *J Immunol.* 1996 Jul 1;157(1):48-56.

*Cherukuri A, Cheng PC, Pierce SK: The role of the CD19/CD21 complex in B cell processing and presentation of complement-tagged antigens. *J Immunol.* 2001 Jul 1;167(1):163-72.

*Krop I, de Fougères AR, Hardy RR, Allison M, Schlissel MS, Fearon DT: Self-renewal of B-1 lymphocytes is dependent on CD19. *Eur J Immunol.* 1996 Jan;26(1):238-42.

*Inabe K, Kurosaki T: Tyrosine phosphorylation of B-cell adaptor for phosphoinositide 3-kinase is required for Akt activation in response to CD19 engagement. *Blood.* 2002 Jan 15;99(2):584-9.

*Shoham T, Rajapaksa R, Boucheix C, Rubinstein E, Poe JC, Tedder TF, Levy S: The tetraspanin CD81 regulates the expression of CD19 during B cell development in a postendoplasmic reticulum compartment. *J Immunol.* 2003 Oct 15;171(8):4062-72.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz