



1P-648-T100

Monoclonal Antibody to CD36 Phycoerythrin (PE) conjugated (100 tests)

Clone:	CB38 (NL07)
Isotype:	Mouse IgM
Specificity:	The mouse monoclonal antibody CB38 (NL07) recognizes CD36 (GPIIb), a 85-113 kDa integral membrane glycoprotein expressed on platelets, macrophages, endothelial cells, early erythroid cells and megakaryocytes. HLDA IV.; WS Code P106
Regulatory Status:	RUO
Immunogen:	living human myeloid cells
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing Tris buffered saline (TBS) solution containing 15 mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD36 (fatty acid translocase, FAT) is an 85-113 kDa ditopic glycosylated protein that belongs to the class B family of scavenger receptors. CD36 is expressed by most resting marginal zone B cells but not by follicular and B1 B cells, and it is rapidly induced on Follicular B cells in vitro upon TLR and CD40 stimulation. CD36 does not affect the development of B cells, but modulates both primary and secondary antibody response. Similarly to glucose transporter GLUT4, CD36 is translocated from intracellular pools to the plasma membrane following cell stimulation by insulin. In mouse, CD36 is responsible for gustatory perception of long-chain fatty acids.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Alessio M, Ghigo D, Garbarino G, Geuna M, Malavasi F: Analysis of the human CD36 leucocyte differentiation antigen by means of the monoclonal antibody NL07. *Cell Immunol.* 1991 Oct 15;137(2):487-500.
- *Alessio M, Roggero S, Bussolino F, Saitta M, Malavasi F: Characterization of the murine monoclonal antibody NL07 specific for the human thrombospondin receptor (CD36 molecule). *Curr Stud Hematol Blood Transfus.* 1991;(58):182-6.
- *Alessio M, Greco NJ, Primo L, Ghigo D, Bosia A, Tandon NN, Ockenhouse CF, Jamieson GA, Malavasi F: Platelet activation and inhibition of malarial cytoadherence by the anti-CD36 IgM monoclonal antibody NL07. *Blood.* 1993 Dec 15;82(12):3637-47.
- *Fadok VA, Warner ML, Bratton DL, Henson PM: CD36 is required for phagocytosis of apoptotic cells by human macrophages that use either a phosphatidylserine receptor or the vitronectin receptor (alpha v beta 3). *J Immunol.* 1998 Dec 1;161(11):6250-7.
- *Matasić R, Dietz AB, Vuk-Pavlović S: Dexamethasone inhibits dendritic cell maturation by redirecting differentiation of a subset of cells. *J Leukoc Biol.* 1999 Dec;66(6):909-14.
- *Hoffmann PR, deCathelineau AM, Ogden CA, Leverrier Y, Bratton DL, Daleke DL, Ridley AJ, Fadok VA, Henson PM: Phosphatidylserine (PS) induces PS receptor-mediated macropinocytosis and promotes clearance of apoptotic cells. *J Cell Biol.* 2001 Nov 12;155(4):649-59.
- *Kapinsky M, Torzewski M, Büchler C, Duong CQ, Rothe G, Schmitz G: Enzymatically degraded LDL preferentially binds to CD14(high) CD16(+) monocytes and induces foam cell formation mediated only in part by the class B scavenger-receptor CD36. *Arterioscler Thromb Vasc Biol.* 2001 Jun;21(6):1004-10.
- *McKenna RW, Washington LT, Aquino DB, Picker LJ, Kroft SH: Immunophenotypic analysis of hematogones (B-lymphocyte precursors) in 662 consecutive bone marrow specimens by 4-color flow cytometry. *Blood.* 2001 Oct 15;98(8):2498-507.
- *Moniuszko M, Kowal K, Rusak M, Pietruczuk M, Dabrowska M, Bodzenta-Lukaszyk A: Monocyte CD163 and CD36 expression in human whole blood and isolated mononuclear cell samples: influence of different anticoagulants. *Clin Vaccine Immunol.* 2006 Jun;13(6):704-7.
- *And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz