



1P-707-T100

Monoclonal Antibody to CD195 / CCR5 Phycoerythrin (PE) conjugated (100 tests)

Clone:	T21/8
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kDa G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.
Regulatory Status:	RUO
Immunogen:	CCR5 peptide (Met1-Lys22) KLH conjugate
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD195 / CCR5 (also known as CKR-5) is a receptor for inflammatory CC-chemokines (characterized by a pair of adjacent cysteine residues), such as MIP-1 alpha, MIP-1 beta, or RANTES. It is a G protein-associated seven-pass transmembrane protein expressed on resting T cells with memory/effector phenotype, monocytes, macrophages and immature dendritic cells. This chemokine receptor regulates the activation and directed migration of leukocytes. Importantly, along with CD4, CD195 / CCR5 functions as a major receptor for HIV. Their ligand is the viral glycoprotein gp120.
References:	*Pollok-Kopp B, Schwarze K, Baradari VK, Oppermann M: Analysis of ligand-stimulated CC chemokine receptor 5 (CCR5) phosphorylation in intact cells using phosphosite-specific antibodies. J Biol Chem. 2003 Jan 24;278(4):2190-8. *Hüttenrauch F, Pollok-Kopp B, Oppermann M: G protein-coupled receptor kinases promote phosphorylation and beta-arrestin-mediated internalization of CCR5 homo- and hetero-oligomers. J Biol Chem. 2005 Nov 11;280(45):37503-15. *Monde K, Maeda Y, Tanaka Y, Harada S, Yusa K: Gp120 V3-dependent impairment of R5 HIV-1 infectivity due to virion-incorporated CCR5. J Biol Chem. 2007 Dec 21;282(51):36923-32. *Mascalchi P, Lamort AS, Salomé L, Dumas F: Single Particle Tracking reveals two distinct environments for CD4 receptors at the surface of living T lymphocytes. Biochem Biophys Res Commun. 2012 Jan 6;417(1):409-13

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz