



1P-755-C025

Monoclonal Antibody to CD266 / TWEAK R Phycoerythrin (PE) conjugated (0.025 mg)

Clone:	ITEM-4
Isotype:	Mouse IgG2b
Specificity:	The mouse monoclonal antibody ITEM-4 recognizes CD266 / TWEAK R, a TNFR superfamily receptor for CD255 / TWEAK, a TNF-like weak inducer of apoptosis.
Regulatory Status:	RUO
Immunogen:	human CD266-transfected P815 cells
Species Reactivity:	Human, Mouse
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD266 / TWEAK R (TNFRSF12A), also known as FN14 (fibroblast growth factor-inducible 14) is a receptor for CD255 / TWEAK, the TNF-like weak inducer of apoptosis. CD266 is expressed on endothelial cells, as well as on some cancer tissues, and plays a role in CD255-induced endothelial cell migration, proliferation, and angiogenesis. The CD255-CD266 interaction, or antibody-mediated triggering of CD266 is also able to induce apoptosis and necrosis in CD266-positive cells (including tumor cells), which might have therapeutic potential.
References:	*Nakayama M, Ishidoh K, Kojima Y, Harada N, Kominami E, Okumura K, Yagita H: Fibroblast growth factor-inducible 14 mediates multiple pathways of TWEAK-induced cell death. <i>J Immunol.</i> 2003 Jan 1;170(1):341-8. *Yoriki R, Akashi S, Sho M, Nomi T, Yamato I, Hotta K, Takayama T, Matsumoto S, Wakatsuki K, Migita K, Yagita H, Nakajima Y: Therapeutic potential of the TWEAK/Fn14 pathway in intractable gastrointestinal cancer. <i>Exp Ther Med.</i> 2011 Jan;2(1):103-108 *Sanz AB, Sanchez-Niño MD, Carrasco S, Manzarbeitia F, Ruiz-Andres O, Selgas R, Ruiz-Ortega M, Gonzalez-Enguita C, Egido J, Ortiz A: Inflammatory cytokines and survival factors from serum modulate tweek-induced apoptosis in PC-3 prostate cancer cells. <i>PLoS One.</i> 2012;7(10):e47440. *Roos C, Wicovsky A, Müller N, Salzmann S, Rosenthal T, Kalthoff H, Trauzold A, Seher A, Henkler F, Kneitz C, Wajant H: Soluble and transmembrane TNF-like weak inducer of apoptosis differentially activate the classical and noncanonical NF-kappa B pathway. <i>J Immunol.</i> 2010 Aug 1;185(3):1593-605.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz