



1P-763-C100

Monoclonal Antibody to CD41 (mouse) Phycoerythrin (PE) conjugated (0.1 mg)

Clone:	MWReg30
Isotype:	Rat IgG1
Specificity:	The rat monoclonal antibody MWReg30 recognizes CD41 (GPIIb), a transmembrane glycoprotein (integrin family) composed of two chains GPIIb alpha (heavy chain; 120 kDa) and GPIIb beta (light chain; 23 kDa). CD41 is mainly expressed on platelets and megakaryocytes.
Regulatory Status:	RUO
Immunogen:	Murine platelets
Species Reactivity:	Mouse
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate was purified by size-exclusion chromatography.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a variety of proteins including von Willebrand factor, fibrinogen, fibronectin and vitronectin. CD41 is mainly expressed on megakaryocyte-platelet lineage, but generally belongs to the antigens that are expressed during early stages of hematopoietic differentiation.
References:	*Brancaleone V, Gobbetti T, Cenac N, le Faouder P, Colom B, Flower RJ, Vergnolle N, Nourshargh S, Perretti M: A vasculo-protective circuit centered on lipoxin A4 and aspirin-triggered 15-epi-lipoxin A4 operative in murine microcirculation. <i>Blood</i> . 2013 Jul 25;122(4):608-17. *Boisset JC, Clapes T, Van Der Linden R, Dzierzak E, Robin C: Integrin αIIb (CD41) plays a role in the maintenance of hematopoietic stem cell activity in the mouse embryonic aorta. <i>Biol Open</i> . 2013 Apr 11;2(5):525-32. *Sullivan BP, Wang R, Tawfik O, Luyendyk JP: Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. <i>Toxicol Sci</i> . 2010 May;115(1):286-94. *Ramsey H, Zhang Q, Brown DE, Steensma DP, Lin CP, Wu MX: Stress-induced hematopoietic failure in the absence of immediate early response gene X-1. <i>Haematologica</i> . 2013 Sep 20. [Epub ahead of print]

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz