



Antibodies

1P-765-T100

Monoclonal Antibody to Perforin Phycoerythrin (PE) conjugated (100 tests)

| | |
|-----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Clone: | dG9 |
| Isotype: | Mouse IgG2b |
| Specificity: | The mouse monoclonal antibody dG9 (also known as deltaG9) recognizes perforin, a 70 kDa protein expressed in cytoplasmic granules of cytotoxic T cells and NK cells. |
| Regulatory Status: | RUO |
| Immunogen: | purified granules from human YT lymphoma cell line |
| Species Reactivity: | Human, Bovine |
| Negative Species: | Mouse |
| Preparation: | The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. |
| Storage Buffer: | The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. |
| Storage / Stability: | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. |
| Usage: | The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |
| Expiration: | See vial label |
| Lot Number: | See vial label |
| Background: | Perforin is a 70 kDa cytolytic protein with structural and functional similarities to complement component 9 (C9). It is stored in cytoplasmic granules of cytotoxic T cells and NK cells and after its release it forms transmembrane pores in the target cells to kill it. As perforin is a key effector molecule for cell-mediated cytotoxicity, defects of its gene can cause severe disorders. |

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Hameed A, Olsen KJ, Cheng L, Fox WM, Hruban RH, Podack ER: Immunohistochemical identification of cytotoxic lymphocytes using human perforin monoclonal antibody. *Am J Pathol.* 1992 May;140(5):1025-30.
 - *Hersperger AR, Makedonas G, Betts MR: Flow cytometric detection of perforin upregulation in human CD8 T cells. *Cytometry A.* 2008 Nov;73(11):1050-7.
 - *Vargas-Inchaustegui DA, Demberg T, Robert-Guroff M: A CD8⁺ subpopulation of macaque circulatory natural killer cells can mediate both antibody-dependent and antibody-independent cytotoxic activities. *Immunology.* 2011 Nov;134(3):326-40.
 - *Takeuchi M, Inoue T, Otani T, Yamasaki F, Nakamura S, Kibata M: Cell-in-cell structures formed between human cancer cell lines and the cytotoxic regulatory T-cell line HOZOT. *J Mol Cell Biol.* 2010 Jun;2(3):139-51.
 - *Qiu Y, Chen J, Liao H, Zhang Y, Wang H, Li S, Luo Y, Fang D, Li G, Zhou B, Shen L, Chen CY, Huang D, Cai J, Cao K, Jiang L, Zeng G, Chen ZW: Tim-3-expressing CD4⁺ and CD8⁺ T cells in human tuberculosis (TB) exhibit polarized effector memory phenotypes and stronger anti-TB effector functions. *PLoS Pathog.* 2012;8(11):e1002984.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz