



Antibodies

1P-778-T025

## Monoclonal Antibody to CD101 Phycoerythrin (PE) conjugated (25 tests)

<b>Clone:</b>	BB27
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody BB27 recognizes CD101, a 140 kDa disulfide-bonded homodimeric protein expressed on activated T cells, and some other cell types, such as granulocytes and cells of the monocyte/macrophage lineage. HLDA V; WS Code T040
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Human thymic clone B12
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD101 is a type I transmembrane glycoprotein, which forms disulfide-linked homodimers. It is expressed on activated T cells, as well as on granulocytes, monocytes, dendritic cells or mucosal T cells. It plays a major role in the activation of T cells by skin dendritic cells. Function of CD101 has not been fully elucidated, but in mice its knock-out results in liver autoimmune disease induced by <i>Novosphingobium aromaticivorans</i> .
<b>References:</b>	*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995). Bagot M, Martinel I, Charue D, Weill F, Boulland ML, Wechsler J, Freeman GJ, Bensussan A, Boumsell L: CD101 is expressed by skin dendritic cells. Role in T-lymphocyte activation. <i>Tissue Antigens</i> . 1997 Nov;50(5):439-48. *Bagot M, Martinel I, Charue D, Boulland ML, Wechsler J, Bensussan A, Boumsell L: Functional role of CD101 on skin dendritic cells. <i>Adv Exp Med Biol</i> . 1997;417:227-32. *Grassi F, Dezutter-Dambuyant C, McIlroy D, Jacquet C, Yoneda K, Imamura S, Boumsell L, Schmitt D, Autran B, Debré P, Hosmalin A: Monocyte-derived dendritic cells have a phenotype comparable to that of dermal dendritic cells and display ultrastructural granules distinct from Birbeck granules. <i>J Leukoc Biol</i> . 1998 Oct;64(4):484-93.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)