



A4-297-T025

Monoclonal Antibody to CD34 Alexa Fluor® 488 conjugated (25 tests)

Clone:	4H11[APG]
Isotype:	Mouse IgG1
Specificity:	The antibody 4H11[APG] reacts with Class III epitope on CD34 (Mucosialin), a 110-115 kDa monomeric transmembrane phosphoglycoprotein expressed on hematopoietic progenitors cells and on the most pluripotential stem cells; it is gradually lost on progenitor cells. The antibody 4H11[APG] completely blocks binding of Class II antibody QBEnd10 and Class III antibodies BIRMA K3 and 8G12 on KG1a cell line. HLDA VI; WS Code M MA58
Regulatory Status:	RUO
Immunogen:	Permanent human cell line derived from peripheral leucocytes of a patient suffering from chronic myeloid leukaemia.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Alexa Fluor® 700 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD34 is a highly glycosylated monomeric 111-115 kDa surface protein, which is present on many stem cell populations. It is a well established stem cell marker, though its expression on human hematopoietic stem cells is reversible. CD34 probably serves as a surface receptor that undergoes receptor-mediated endocytosis and regulates adhesion, differentiation and proliferation of hematopoietic stem cells and other progenitors. CD34 expression is likely to represent a specific state of hematopoietic development that may have altered adhering properties with expanding and differentiating capabilities in both in vitro and in vivo conditions.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Krauter J, Hartl M, Hambach L, Kohlenberg A, Gunsilius E, Ganser A, Heil G: Receptor-mediated endocytosis of CD34 on hematopoietic cells after stimulation with the monoclonal antibody anti-HPCA-1. *J Hematother Stem Cell Res.* 2001 Dec;10(6):863-71.
- *Dao MA, Arevalo J, Nolte JA: Reversibility of CD34 expression on human hematopoietic stem cells that retain the capacity for secondary reconstitution. *Blood.* 2003 Jan 1;101(1):112-8.
- *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Gulati S, Luthra PM: Three-dimensional structure prediction of the interaction of CD34 with the SH3 domain of Crk-L. *Stem Cells Dev.* 2005 Oct;14(5):470-7.
- *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Luthra PM: Hematopoietic stem cell antigen CD34: role in adhesion or homing. *Stem Cells Dev.* 2006 Jun;15(3):305-13.
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- *Elknerová K, Lacinová Z, Soucek J, Marinov I, Stöckbauer P: Growth inhibitory effect of the antibody to hematopoietic stem cell antigen CD34 in leukemic cell lines. *Neoplasma.* 2007;54(4):311-20.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.