

A4-309-T100

Monoclonal Antibody to CD41 Alexa Fluor® 488 conjugated (100 tests)

Clone: MEM-06

Isotype: Mouse IgG1

Specificity: The antibody MEM-06 reacts with CD41 (GPIIb), a transmembrane glycoprotein

(integrin family) composed of two chains GPIIb alpha (heavy chain; 120 kDa) and GPIIb beta (light chain; 23 kDa). CD41 is mainly expressed on platelets and

megakaryocytes. Workshop: HLDA 10

Regulatory Status: RUO

Immunogen: Leukocytes of patient suffering from LGL-type leukaemia.

Species Reactivity: Human

Preparation: The purified antibody is conjugated with Alexa Fluor® 488 under optimum

conditions. The conjugate is purified by size-exclusion chromatography and

adjusted for direct use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in stabilizing phosphate buffered saline (PBS) solution

containing 15mM sodium azide.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not

use after expiration date stamped on vial label.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 4

μl reagent / 100 μl of whole blood or 10⁶ cells in a suspension.

The content of a vial (0.4 ml) is sufficient for 100 tests.

Expiration: See vial label

Lot Number: See vial label

Background: CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and

23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a variety of proteins including von Willebrand factor, fibrinogen, fibronectin and vitronectin. CD41 is mainly expressed on megakaryocyte-platelet lineage, but generally belongs to the antigens that are expressed during early stages of

hematopoietic differentiation.



PRODUCT DATA SHEET

References:

*Debili N, Robin C, Schiavon V, Letestu R, Pflumio F, Mitjavila-Garcia MT, Coulombel L, Vainchenker W: Different expression of CD41 on human lymphoid and myeloid progenitors from adults and neonates. Blood. 2001 Apr 1;97(7):2023-30.

*Mitjavila-Garcia MT, Cailleret M, Godin I, Nogueira MM, Cohen-Solal K, Schiavon V, Lecluse Y, Le Pesteur F, Lagrue AH, Vainchenker W: Expression of CD41 on hematopoietic progenitors derived from embryonic hematopoietic cells. Development. 2002 Apr;129(8):2003-13.

*Ferkowicz MJ, Starr M, Xie X, Li W, Johnson SA, Shelley WC, Morrison PR, Yoder MC: CD41 expression defines the onset of primitive and definitive hematopoiesis in the murine embryo. Development. 2003 Sep;130(18):4393-403.

*Zhang J, Varas F, Stadtfeld M, Heck S, Faust N, Graf T: CD41-YFP mice allow in vivo labeling of megakaryocytic cells and reveal a subset of platelets hyperreactive to thrombin stimulation. Exp Hematol. 2007 Mar;35(3):490-499.

*Pagani G, Pereira JPV, Stoldt VR, Beck A, Scharf RE, Gohlke H: The human platelet antigen-1b (Pro33) variant of αIlbβ3 allosterically shifts the dynamic conformational equilibrium of this integrin toward the active state. J Biol Chem. 2018 Mar 30;293(13):4830-4844.

*Belizaire RM, Prakash PS, Richter JR, Robinson BR, Edwards MJ, Caldwell CC, Lentsch AB, Pritts TA: Microparticles from stored red blood cells activate neutrophils and cause lung injury after hemorrhage and resuscitation. J Am Coll Surg. 2012 Apr;214(4):648-55

*Komsa-Penkova R, Todinova SJ, Andreeva TD, Krumova SB, Taneva SG, Golemanov GM, Georgieva GA, Mihaylova NM, Tchorbanov PT: Alterations in platelet activity and elastic modulus of healthy subjects, carriers of G20210A polymorphism in the prothrombin gene. J Biomed Clin Res Volume 9 Number 1, 2016

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.