

A6-224-T100

## Monoclonal Antibody to CD45RB Alexa Fluor® 647 conjugated (100 tests)

<b>Clone:</b>	MEM-55
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	<p>The antibody MEM-55 recognizes a sialidase-sensitive epitope of CD45RB, a 180-240 kDa single chain type I membrane glycoprotein, variant of CD45 (CD45RB isoform). CD45RB is expressed on a subset of T lymphocytes, B lymphocytes, monocytes, macrophages, granulocytes and dendritic cells.</p> <p>HLDA III; WS Code NL 358 HLDA IV; WS Code NL 2 HLDA V; WS Code T T-151 HLDA V; WS Code T T-CD45.08</p>
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Human thymocytes and T lymphocytes.
<b>Species Reactivity:</b>	Human, Non-Human Primates
<b>Preparation:</b>	The purified antibody is conjugated with Alexa Fluor® 647 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension.</p> <p>The content of a vial (0.4 ml) is sufficient for 100 tests.</p>
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<p>CD45RB is an of a receptor-type protein tyrosine phosphatase, CD45 glycoprotein. CD45 is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases, promotes cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis. CD45 isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. CD45RB is expressed e.g. in microglia and inflammatory cells.</p>

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Townsend KP, Vendrame M, Ehrhart J, Faza B, Zeng J, Town T, Tan J: CD45 isoform RB as a molecular target to oppose lipopolysaccharide-induced microglial activation in mice. *Neurosci Lett*. 2004 May 13;362(1):26-30.
- \*Li FJ, Tsuyama N, Ishikawa H, Obata M, Abroun S, Liu S, Otsuyama K, Zheng X, Ma Z, Maki Y, Kawano MM: A rapid translocation of CD45RO but not CD45RA to lipid rafts in IL-6-induced proliferation in myeloma. *Blood*. 2005 Apr 15;105(8):3295-302.
- \*Cosenza-Nashat MA, Kim MO, Zhao ML, Suh HS, Lee SC: CD45 isoform expression in microglia and inflammatory cells in HIV-1 encephalitis. *Brain Pathol*. 2006 Oct;16(4):256-65.
- \*Dawes R, Petrova S, Liu Z, Wraith D, Beverley PC, Tchilian EZ. Combinations of CD45 isoforms are crucial for immune function and disease. *J Immunol*. 2006 Mar 15;176(6):3417-25.
- \*Bijian K, Zhang L, Shen SH: Collagen-mediated survival signaling is modulated by CD45 in Jurkat T cells. *Mol Immunol*. 2007 Jul;44(15):3682-90.
- \*Desharnais P, Dupéré-Minier G, Hamelin C, Devine P, Bernier J: Involvement of CD45 in DNA fragmentation in apoptosis induced by mitochondrial perturbing agents. *Apoptosis*. 2007 Dec 19
- \*Leukocyte Typing III., McMichael A. J. et al (Eds.), Oxford University Press (1987).
- \*Horejsi V, Angelisova P, Bazil V, Kristofova H, Stoyanov S, Stefanova I, Hausner P, Vosecky M, Hilgert I: Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). *Folia Biol (Praha)*. 1988;34(1):23-34.
- \*Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- \*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- \*Koethe S, Zander L, Köster S, Annan A, Ebenfelt A, Spencer J, Bemark M: Pivotal advance: CD45RB glycosylation is specifically regulated during human peripheral B cell differentiation. *J Leukoc Biol*. 2011 Jul;90(1):5-19.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)