



A7-342-T025

## Monoclonal Antibody to CD46 Alexa Fluor® 700 conjugated (25 tests)

<b>Clone:</b>	MEM-258
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The antibody MEM-258 recognizes an epitope on SCR4 (the membrane-proximal SCR) domain of CD46 (Membrane cofactor protein). CD46 is 56-66 kDa dimeric transmembrane protein expressed on T and B lymphocytes, platelets, monocytes, granulocytes, endothelial cells, epithelial cells and fibroblast; it is negative on erythrocytes.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	HPB-ALL human T cell line
<b>Species Reactivity:</b>	Human, Bovine
<b>Preparation:</b>	The purified antibody is conjugated with Alexa Fluor® 700 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD46 (MCP, membrane cofactor protein) is a multifunctional cell surface transmembrane protein that binds and inactivates C3b and C4b complement fragments, regulates T cell-induced inflammatory responses by either inhibiting (CD46-1 isoform) or increasing (CD46-2 isoform) the contact hypersensitivity reaction. CD46 also serves as a receptor for several human pathogens (both bacteria and viruses), and its ligation alters T lymphocyte polarization toward antigen-presenting cells or target cells, inhibiting lymphocyte function. CD46 is a protector of placental tissue and is also expressed on the inner acrosomal membrane of spermatozoa.

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Okada N, Liszewski MK, Atkinson JP, Caparon M: Membrane cofactor protein (CD46) is a keratinocyte receptor for the M protein of the group A streptococcus. *Proc Natl Acad Sci U S A*. 1995 Mar 28;92(7):2489-93.
- \*Marie JC, Astier AL, Rivallier P, Rabourdin-Combe C, Wild TF, Horvat B: Linking innate and acquired immunity: divergent role of CD46 cytoplasmic domains in T cell induced inflammation. *Nat Immunol*. 2002 Jul;3(7):659-66.
- \*Gaggar A, Shayakhmetov DM, Lieber A: CD46 is a cellular receptor for group B adenoviruses. *Nat Med*. 2003 Nov;9(11):1408-12.
- \*Liszewski MK, Kemper C, Price JD, Atkinson JP: Emerging roles and new functions of CD46. *Springer Semin Immunopathol*. 2005 Nov;27(3):345-58.
- \*Oliaro J, Pasam A, Waterhouse NJ, Browne KA, Ludford-Menting MJ, Trapani JA, Russell SM: Ligation of the cell surface receptor, CD46, alters T cell polarity and response to antigen presentation. *Proc Natl Acad Sci U S A*. 2006 Dec 5;103(49):18685-90.
- \*Fleischli C, Verhaagh S, Havenga M, Sirena D, Schaffner W, Cattaneo R, Greber UF, Hemmi S: The distal short consensus repeats 1 and 2 of the membrane cofactor protein CD46 and their distance from the cell membrane determine productive entry of species B adenovirus serotype 35. *J Virol*. 2005 Aug;79(15):10013-22.
- \*Weyand NJ, Lee SW, Higashi DL, Cawley D, Yoshihara P, So M: Monoclonal antibody detection of CD46 clustering beneath *Neisseria gonorrhoeae* microcolonies. *Infect Immun*. 2006 Apr;74(4):2428-35.
- \*Fremeaux-Bacchi V, Moulton EA, Kavanagh D, Dragon-Durey MA, Blouin J, Caudy A, Arzouk N, Cleper R, Francois M, Guest G, Pourrat J, Seligman R, Fridman WH, Loirat C, Atkinson JP: Genetic and functional analyses of membrane cofactor protein (CD46) mutations in atypical hemolytic uremic syndrome. *J Am Soc Nephrol*. 2006 Jul;17(7):2017-25.
- \*Loré K, Adams WC, Havenga MJ, Precopio ML, Holterman L, Goudsmit J, Koup RA: Myeloid and plasmacytoid dendritic cells are susceptible to recombinant adenovirus vectors and stimulate polyfunctional memory T cell responses. *J Immunol*. 2007 Aug 1;179(3):1721-9.
- \*Fleischli C, Sirena D, Lesage G, Havenga MJ, Cattaneo R, Greber UF, Hemmi S: Species B adenovirus serotypes 3, 7, 11 and 35 share similar binding sites on the membrane cofactor protein CD46 receptor. *J Gen Virol*. 2007 Nov;88(Pt 11):2925-34.
- \*Hoffmann D, Bayer W, Heim A, Potthoff A, Nettelbeck DM, Wildner O: Evaluation of twenty-one human adenovirus types and one infectivity-enhanced adenovirus for the treatment of malignant melanoma. *J Invest Dermatol*. 2008 Apr;128(4):988-98.
- \*Wang H, Tuve S, Erdman DD, Lieber A: Receptor usage of a newly emergent adenovirus type 14. *Virology*. 2009 May 10;387(2):436-41.
- \*Rebetz J, Na M, Su C, Holmqvist B, Edqvist A, Nyberg C, Widegren B, Salford LG, Sjögren HO, Arnberg N, Qian Q, Fan X: Fiber mediated receptor masking in non-infected bystander cells restricts adenovirus cell killing effect but promotes adenovirus host co-existence. *PLoS One*. 2009 Dec 29;4(12):e8484.
- \*Kälin S, Amstutz B, Gastaldelli M, Wolfrum N, Boucke K, Havenga M, DiGennaro F, Liska N, Hemmi S, Greber UF: Macropinocytotic uptake and infection of human epithelial cells with species B2 adenovirus type 35. *J Virol*. 2010 May;84(10):5336-50.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)