

PB-506-T100

Monoclonal Antibody to CD22 Pacific Blue™ conjugated (100 tests)

Clone: IS7

Isotype: Mouse IgG1

Specificity: The antibody IS7 reacts with CD22 (BL-CAM), a 130 kDa type I transmembrane

glycoprotein (immunoglobulin superfamily) expressed in the cytoplasm of pro-B and pre-B lymphocytes, and on the surface of mature and activated B lymphocytes; it is lost on plasma cells, peripheral blood T lymphocytes,

granulocytes and monocytes. HLDA IV; WS Code B 227 HLDA V; WS Code B CD22.8

Regulatory Status: RUO

Immunogen: human cell line Reh

Species Reactivity: Human

Preparation: The purified antibody is conjugated with Pacific Blue™ under optimum conditions.

The conjugate is purified by size-exclusion chromatography and adjusted for direct

use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in stabilizing phosphate buffered saline (PBS) solution

containing 15mM sodium azide.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not

use after expiration date stamped on vial label.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 4

μl reagent / 100 μl of whole blood or 10⁶ cells in a suspension.

The content of a vial (0.4 ml) is sufficient for 100 tests.

Expiration: See vial label

Lot Number: See vial label

Background:

CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through

the immunoreceptor tyrosine-based inhibitory motifs (ITIMs), thus setting a treshold for B cell receptor-mediated activation. CD22 also regulates B-cell response by involvement in controlling the CD19/CD21-Src-family protein tyrosine kinase amplification pathway and CD40 signaling. CD22 exhibits hallmarks of

clathrin-mediated endocytic pathway.

References: *Tedder TF, Poe JC, Haas KM: CD22: A Multifunctional Receptor That Regulates

B Lymphocyte Survival and Signal Transduction. Adv Immunol. 2005;88:1-50. *Tateno H, Li H, Schur MJ, Bovin N, Crocker PR, Wakarchuk WW, Paulson JC: Distinct endocytic mechanisms of CD22 (Siglec-2) and Siglec-F reflect roles in cell

signaling and innate immunity. Mol Cell Biol. 2007 Aug;27(16):5699-710.

*Walker JA, Smith KG: CD22: an inhibitory enigma. Immunology. 2007 Dec 7

*Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).

*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).

For laboratory research only, not for drug, diagnostic or other use.



PRODUCT DATA SHEET

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.