



PC-213-T025

## Monoclonal Antibody to CD15 PerCP (25 tests)

<b>Clone:</b>	MEM-158
<b>Isotype:</b>	Mouse IgM
<b>Specificity:</b>	The antibody MEM-158 reacts with CD15, a cell membrane molecule 3-fucosyl-N-acetyllactosamine (3-FAL) strongly expressed on granulocytes, monocytes, macrophages, mast cells; it is also present on Langerhans cells and some myeloid precursors cells. HLDA VI; WS Code AS A053
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Human granulocytes
<b>Species Reactivity:</b>	Human
<b>Negative Species:</b>	Porcine, Bovine, Sheep
<b>Preparation:</b>	The purified antibody is conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing Tris buffered saline (TBS) solution containing 15 mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD15 (Lewis X, Le(x); stage specific embryonic antigen-1, SSEA-1) is a trisaccharide determinant (3-fucosyl-N-acetyllactosamine) expressed on several glycolipids, glycoproteins and proteoglycans of various cell types, e.g. granulocytes, mast cells, monocytes, macrophages, cells of gastric mucosa, nervous system or various tumour cells. There are several variants of Lewis x, such as sialyl-Lewis x or sulphated Lewis x. Cells with high surface expression of Le(x) antigen exhibit strong self-aggregation, based on calcium-dependent Le(x)-Le(x) interaction. This process is involved for example in embryo compaction or in autoaggregation of teratocarcinoma cells. Sialyl-Le(x) and its isomer sialyl-Le(a) are ligands of selectins. CD15 expression has been extensively used to confirm diagnosis of Hodgkin's disease.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*Benharroch D, Dima E, Levy A, Ohana-Malka O, Ariad S, Prinsloo I, Mejirovsky E, Sacks M, Gopas J: Differential expression of sialyl and non-sialyl-CD15 antigens on Hodgkin-Reed-Sternberg cells: significance in Hodgkin's disease. *Leuk Lymphoma*. 2000 Sep;39(1-2):185-94.
  - \*Hakomori S: Le(X) and related structures as adhesion molecules. *Histochem J*. 1992 Nov;24(11):771-6.
  - \*Li C, Wong P, Pan T, Xiao F, Yin S, Chang B, Kang SC, Ironside J, Sy MS: Normal cellular prion protein is a ligand of selectins: binding requires Le(X) but is inhibited by sLe(X). *Biochem J*. 2007 Sep 1;406(2):333-41.
  - \*Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)