

PC-474-T025

Monoclonal Antibody to HLA-DR PerCP (25 tests)

Clone: **MEM-12** Isotype: Mouse IgG1 **Specificity:** The antibody MEM-12 recognizes common epitope on human HLA-DR which is dependent on the association of alpha and beta chains. DR is the isotype of human MHC Class II molecules expressed on antigen-presenting cells (APC; dendritic cells, B lymphocytes, monocytes, macrophages). **Regulatory Status:** RUO thymocyte membrane Immunogen: **Species Reactivity:** Human **Preparation:** The purified antibody is conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. Storage Buffer: The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 10° cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests. **Expiration:** See vial label Lot Number: See vial label **Background:** HLA-DR, a member of MHC class II glycoproteins, that bind intracellularly processed peptides and present them to the Th cells, is composed of 36 kDa alpha chain and 27 kDa beta chain, both anchored in the plasma membrane. Together with other MHC II molecules HLA-DR plays a central role in the immune system. **References:** *Horejsi V, Nemec M, Angelisova P, Kristofova H, Gorga JC, Hilgert I: Characterization of seven new monoclonal antibodies against human DR. DR + DP and DQ1 + DQ3 antigens. Tissue Antigens. 1986 Nov;28(5):288-97.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz