



T7-700-T025

## Monoclonal Antibody to CD123 PE-Cy™7 conjugated (25 tests)

<b>Clone:</b>	6H6
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody 6H6 recognizes CD123 (interleukin 3 receptor alpha), a 60-70 kDa transmembrane protein expressed by myeloid precursors, megakaryocytes, macrophages, dendritic cells, mast cells, basophils, and some B cells. This antibody does not inhibit IL-3 binding to its receptor.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	IL3 receptor alpha chain expressed on the surface of transiently transfected COS cells
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with tandem dye PE-Cy™7 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD123 is the alpha chain of interleukin 3 receptor (IL-3R alpha). This subunit heterodimerizes with the interleukin 3 receptor beta chain (CD131), which is shared with other receptors. CD123 interacts with IL-3 specifically, but with low affinity, and association with the beta subunit confers high affinity binding to the receptor heterodimer. Both chains are required for signaling, but receptor activation and signal transduction depend on IL-3 binding to CD123 as the initial step.
<b>References:</b>	*Sun Q, Woodcock JM, Rapoport A, Stomski FC, Korpelainen EI, Bagley CJ, Goodall GJ, Smith WB, Gamble JR, Vadas MA, Lopez AF: Monoclonal antibody 7G3 recognizes the N-terminal domain of the human interleukin-3 (IL-3) receptor alpha-chain and functions as a specific IL-3 receptor antagonist. Blood. 1996 Jan 1;87(1):83-92. *Martín-Gayo E, Sierra-Filardi E, Corbí AL, Toribio ML: Plasmacytoid dendritic cells resident in human thymus drive natural Treg cell development. Blood. 2010 Jul 1;115(26):5366-75. *Herling M, Teitell MA, Shen RR, Medeiros LJ, Jones D: TCL1 expression in plasmacytoid dendritic cells (DC2s) and the related CD4+ CD56+ blastic tumors of skin. Blood. 2003 Jun 15;101(12):5007-9.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

Cy™ and CyDye™ are registered trademarks of GE Healthcare.

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)