



T8-319-T025

Monoclonal Antibody to IgG (Fab) PE-Cy™5 conjugated (25 tests)

Clone:	4A11
Isotype:	Mouse IgG2b
Specificity:	The antibody 4A11 reacts with Fab fragment of human IgG.
Regulatory Status:	RUO
Immunogen:	Purified human IgG.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with tandem dye PE-Cy™5 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins (T1/2 = 21 days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.
References:	*Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975;194:77-95. *Fuller JM, Keyser JW: Serum immunoglobulins after surgical operation. Clin Chem. 1975 May;21(6):667-71. *Balogh Z, Merétey K, Falus A, Bozsóky S: Serological abnormalities in juvenile chronic arthritis: a review of 46 cases. Ann Rheum Dis. 1980 Apr;39(2):129-34. *Brinkmann V, Heusser CH: T cell-dependent differentiation of human B cells into IgM, IgG, IgA, or IgE plasma cells: high rate of antibody production by IgE plasma cells, but limited clonal expansion of IgE precursors. Cell Immunol. 1993 Dec;152(2):323-32.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

Cy™ and CyDye™ are registered trademarks of GE Healthcare.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz