



### GTX300058 EED-EZH2 / PRC2 complex Antibody Panel

#### Content

Cat No	Product Name	Applications	Package
GTX112268	EED antibody	ICC/IF, IHC-P, WB	25 µl
GTX110384	EZH2 antibody	ICC/IF, WB, ChIP assay	25 µl
GTX109642	HDAC2 antibody	ICC/IF, IHC-P, IP, WB, ChIP assay	25 µl
GTX100513	HDAC1 antibody	ICC/IF, IHC-P, IP, WB, ChIP assay	25 µl
GTX213110-01	Rabbit IgG antibody (HRP)	Dot, ELISA, IHC-P, WB	25 µl

#### Note

For *In vitro* laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### DataSheet - 1

<b>Catalog Number</b>	GTX112268	Package:25 µl, 100 µl
<b>Product Name</b>	EED antibody	
<b>Full Name</b>	embryonic ectoderm development	
<b>Synonyms</b>	HEED antibody, WAIT1 antibody, EED antibody, WD protein associating with integrin cytoplasmic tails 1 antibody, polycomb protein EED antibody, embryonic ectoderm development antibody	
<b>Product Description</b>	Rabbit Polyclonal antibody to EED (embryonic ectoderm development)	
<b>Background</b>	This gene encodes a member of the Polycomb-group (PcG) family. PcG family members form multimeric protein complexes, which are involved in maintaining the transcriptional repressive state of genes over successive cell generations. This protein interacts with enhancer of zeste 2, the cytoplasmic tail of integrin beta7, immunodeficiency virus type 1 (HIV-1) MA protein, and histone deacetylase proteins. This protein mediates repression of gene activity through histone deacetylation, and may act as a specific regulator of integrin function. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq]	
<b>Host</b>	Rabbit	
<b>Clonality</b>	Polyclonal	
<b>Isotype</b>	IgG	
<b>Immunogen</b>	Recombinant protein encompassing a sequence within the center region of human EED. The exact sequence is proprietary.	
<b>Antigen Species</b>	Human	
<b>Species Reactivity</b>	Human	
<b>Predicted Cross Reactivity species</b>	Mouse, Rat, Zebrafish, Japanese medaka, Chicken, Rhesus Monkey, Bovine	
<b>Predict Reactivity Note</b>	Mouse (100%), Rat (100%), Zebrafish (93%), Japanese medaka (95%), Chicken (99%), Rhesus Monkey (100%), Bovine (100%)	
<b>Applications</b>	ICC/IF, IHC-P, WB	
		<b>Suggested dilution</b>
		<b>Reference</b>
<b>Application Note</b>	ICC/IF	1:100-1:1000*
	IHC (Formalin-fixed paraffin-embedded sections)	1:100-1:1000*

Western blot 1:500-1:3000\*

Not tested in other applications.

\*Optimal dilutions/concentrations should be determined by the researcher.

Positive Controls H1299 , HCT116

Predicted Target Size 50 kDa

Cellular Localization Nucleus

Conjugation Unconjugated

Form Supplied Liquid

Purification Purified by antigen-affinity chromatography.


Concentration 1 mg/ml (Please refer to the vial label for the specific concentration)

Storage Buffer 1XPBS, 20% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.

Storage Instruction Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Notes For *In vitro* laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

ResearchArea [Epigenetics](#) > [Histone acetylation](#) > [Histone deacetylase](#)



#### GTX112268 WB Image


EED antibody detects EED protein by Western blot analysis.

A. 30 µg H1299 whole cell lysate/extract

B. 30 µg HCT116 whole cell lysate/extract


10 % SDS-PAGE

EED antibody (GTX112268) dilution: 1:1000



#### GTX112268 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed A431, using EED(GTX112268) antibody at 1:500 dilution.




#### GTX112268 IHC-P Image

Immunohistochemical analysis of paraffin-embedded SAS Xenograft, using EED(GTX112268) antibody at 1:100 dilution.

#### DataSheet - 2

<b>Catalog Number</b>	GTX110384	Package:25 µl, 100 µl	
<b>Product Name</b>	EZH2 antibody		
<b>Full Name</b>	enhancer of zeste homolog 2 (Drosophila)		
<b>Synonyms</b>	ENX-1 antibody, ENX1 antibody, EZH1 antibody, KMT6 antibody, KMT6A antibody, MGC9169 antibody, EZH2 antibody, histone-lysine N-methyltransferase EZH2 antibody, enhancer of zeste 2 antibody, lysine N-methyltransferase 6 antibody, enhancer of zeste homolog 2 (Drosophila) antibody		
<b>Product Description</b>	Rabbit Polyclonal antibody to EZH2 (histone-lysine N-methyltransferase EZH2)		
<b>Background</b>	This gene encodes a member of the Polycomb-group (PcG) family. PcG family members form multimeric protein complexes, which are involved in maintaining the transcriptional repressive state of genes over successive cell generations. This protein associates with the embryonic ectoderm development protein, the VAV1 oncoprotein, and the X-linked nuclear protein. This protein may play a role in the hematopoietic and central nervous systems. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq]		
<b>Host</b>	Rabbit		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Immunogen</b>	Recombinant protein encompassing a sequence within the center region of human EZH2. The exact sequence is proprietary.		
<b>Antigen Species</b>	Human		
<b>Species Reactivity</b>	Human		
<b>Predicted Cross Reactivity species</b>	Mouse, Rat, Zebrafish, Japanese medaka, Pig, Rhesus Monkey, Bovine		
<b>Predict Reactivity Note</b>	Mouse (99%), Rat (99%), Zebrafish (94%), Japanese medaka (95%), Pig (99%), Rhesus Monkey (100%), Bovine (99%)		
<b>Applications</b>	ICC/IF, WB, ChIP assay		
<b>Application Note</b>		Suggested dilution	Reference
	ChIP assay	Assay-dependent dilution	
	ICC/IF	1:100-1:1000*	
	Western blot	1:500-1:3000*	
	Not tested in other applications. *Optimal dilutions/concentrations should be determined by the researcher.		
<b>Positive Controls</b>	H1299 , HCT116 , MCF-7		
<b>Predicted Target Size</b>	85 kDa		

Conjugation	Unconjugated
Form Supplied	Liquid
Purification	Purified by antigen-affinity chromatography.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration)
Storage Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.
Storage Instruction	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Notes	For <i>In vitro</i> laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
ResearchArea	<a href="#">Cancer</a> > <a href="#">Tumor biomarkers</a> <a href="#">Cancer</a> > <a href="#">Type of cancer</a> > <a href="#">Prostate cancer</a> <a href="#">Epigenetics</a> > <a href="#">Histone acetylation</a> > <a href="#">Histone deacetylase</a>



#### GTX110384 ICC/IF Image

EZH2 antibody detects EZH2 protein at nucleus by immunofluorescent analysis.


Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: EZH2 protein stained by EZH2 antibody (GTX110384) diluted at 1:500.

Red: phalloidin, a cytoskeleton marker, diluted at 1:200.


Blue: Hoechst 33342 staining.

Scale bar = 10 µm.



#### GTX110384 ChIP Image

Cross-linked ChIP was performed with HeLa chromatin extract and 5 µg of either control rabbit IgG or anti-EZH2 antibody. The precipitated DNA was detected by PCR with primer set targeting to CCND2.



**GTX110384 WB Image**

Sample (30 ug of whole cell lysate)

A: H1299

B: HCT116

C: MCF-7


7.5% SDS PAGE

GTX110384 diluted at 1:1000

**DataSheet - 3**


<b>Catalog Number</b>	GTX109642	Package:25 µl, 100 µl	★★★★★ (1)	<a href="#">Reference</a> (1)
<b>Product Name</b>	HDAC2 antibody			
<b>Full Name</b>	histone deacetylase 2			
<b>Synonyms</b>	HD2 antibody, RPD3 antibody, YAF1 antibody, HDAC2 antibody, histone deacetylase 2 antibody, transcriptional regulator homolog RPD3 antibody, YY1-associated factor 1 antibody			
<b>Product Description</b>	Rabbit Polyclonal antibody to HDAC2 (histone deacetylase 2)			
<b>Background</b>	This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes and are responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). This protein also forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus it plays an important role in transcriptional regulation, cell cycle progression and developmental events. [provided by RefSeq]			
<b>Host</b>	Rabbit			
<b>Clonality</b>	Polyclonal			
<b>Isotype</b>	IgG			
<b>Immunogen</b>	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human HDAC2. The exact sequence is proprietary.			
<b>Antigen Species</b>	Human			
<b>Species Reactivity</b>	Human, Mouse, Rat			
<b>Applications</b>	ICC/IF, IHC-P, IP, WB, ChIP assay			
		<b>Suggested dilution</b>	<b>Reference</b>	
	ChIP assay	Assay-dependent dilution		
	ICC/IF	1:100-1:1000*		
<b>Application Note</b>	IHC (Formalin-fixed paraffin-embedded sections)	1:100-1:1000*		
	Immunoprecipitation	1:100-1:500*		
	Western blot	1:500-1:3000*		
	Not tested in other applications.			
	*Optimal dilutions/concentrations should be determined by the researcher.			
<b>Positive Controls</b>	293T , A431 , HeLa , HepG2 , C2C12 , Rat2 , *NB2a			

<b>Predicted Target Size</b>	55 kDa
<b>Cellular Localization</b>	Nucleus
<b>Conjugation</b>	Unconjugated
<b>Form Supplied</b>	Liquid
<b>Purification</b>	Purified by antigen-affinity chromatography.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration)
<b>Storage Buffer</b>	0.1M Tris, 0.1M Glycine, 10% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.
<b>Storage Instruction</b>	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Notes</b>	For <i>In vitro</i> laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
<b>ResearchArea</b>	<a href="#">Cancer</a> > <a href="#">Type of cancer</a> > <a href="#">Breast</a> > <a href="#">Other</a> <a href="#">Cancer</a> > <a href="#">Type of cancer</a> > <a href="#">Ovarian</a> <a href="#">Epigenetics</a> > <a href="#">Histone acetylation</a> > <a href="#">Histone deacetylase</a>




#### GTX109642 ICC/IF Image

Confocal immunofluorescence analysis (Olympus FV10i) of paraformaldehyde-fixed HeLa, using HDAC2(GTX109642) antibody (Green) at 1:500 dilution. Alpha-tubulin filaments were labeled with GTX11304 (Red) at 1:500.



#### GTX109642 WB Image

HDAC2 antibody detects HDAC2 protein by western blot analysis. Various whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with HDAC2 antibody (GTX109642) diluted by 1:1000.




**GTX109642 ChIP Image**

HDAC2 antibody immunoprecipitates HDAC2 protein-DNA complex in ChIP experiments. ChIP Sample: 293T whole cell lysate/extract

A. 5 µg preimmune rabbit IgG


B. 5 µg of HDAC2 antibody (GTX109642)

The precipitated DNA was detected by PCR with primer set targeting to GAPDH promoter.




**GTX109642 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded Cal27 Xenograft, using HDAC2(GTX109642) antibody at 1:100 dilution.



**GTX109642 IP Image**

HDAC2 antibody immunoprecipitates HDAC2 protein in IP experiments. IP Sample: HeLa whole cell lysate/extract A. 40 µg HeLa whole cell lysate/extract B. Control with 2 of preimmune rabbit IgG C. Immunoprecipitation of HDAC2 protein by 2 µg of HDAC2 antibody (GTX109642) 7.5% SDS-PAGE The immunoprecipitated HDAC2 protein was detected by HDAC2 antibody (GTX109642) diluted at 1:1000. EasyBlot anti-rabbit IgG (GTX221666-01) was used as a secondary reagent.




#### GTX109642 WB Image

HDAC2 antibody detects HDAC2 protein by Western blot analysis.

A. 30 µg C2C12 whole cell lysate/extract

7.5 % SDS-PAGE

HDAC2 antibody (GTX109642) dilution: 1:1000



#### GTX109642 WB Image

HDAC2 antibody detects HDAC2 protein by Western blot analysis.

A. 30 µg Rat2 whole cell lysate/extract

7.5 % SDS-PAGE

HDAC2 antibody (GTX109642) dilution: 1:1000

#### Application Reference

1. Lee KH (2014) *Sci Rep* 6394

#### DataSheet - 4

Catalog Number	GTX100513	Package:25 µl, 100 µl	★★★★☆ (1)	<a href="#">Reference</a> (5)
Product Name	HDAC1 antibody			
Full Name	histone deacetylase 1			
Synonyms	DKFZp686H12203 antibody, GON-10 antibody, HD1 antibody, RPD3 antibody, RPD3L1 antibody, HDAC1 antibody, histone deacetylase 1 antibody, reduced potassium dependency, yeast homolog-like 1 antibody			
Product Description	Rabbit Polyclonal antibody to HDAC1 (histone deacetylase 1)			
Background	Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis. [provided by RefSeq]			

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Immunogen</b>	Recombinant protein encompassing a sequence within the center region of human HDAC1. The exact sequence is proprietary.
<b>Antigen Species</b>	Human
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Predicted Cross Reactivity species</b>	Bovine
<b>Predict Reactivity Note</b>	Bovine (98%)
<b>Applications</b>	ICC/IF, IHC-P, IP, WB, ChIP assay

	Suggested dilution	Reference
ChIP assay	Assay-dependent dilution	
ICC/IF	1:100-1:1000*	
IHC (Formalin-fixed paraffin-embedded sections)	1:100-1:1000*	
Immunoprecipitation	1:100-1:500*	
Western blot	1:500-1:3000*	

Not tested in other applications.

\*Optimal dilutions/concentrations should be determined by the researcher.

**Positive Controls** 293T , A431 , HeLa , HepG2 , NIH-3T3 , U87-MG , SK-N-SH , \*E8.5 embryo

**Predicted Target Size** 55 kDa

**Cellular Localization** Nucleus

**Conjugation** Unconjugated

**Form Supplied** Liquid

**Purification** Purified by antigen-affinity chromatography.


**Concentration** 1 mg/ml (Please refer to the vial label for the specific concentration)

**Storage Buffer** 1XPBS, 20% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.

**Storage Instruction** Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

**Notes** For *In vitro* laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**ResearchArea** [Cancer > Apoptosis > Anti-apoptosis](#)  
[Cancer > Type of cancer > Ovarian](#)  
[Cell Biology > Apoptosis > Anti-apoptosis](#)



**GTX100513 ICC/IF Image**


HDAC1 antibody detects HDAC1 protein at nucleus by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: HDAC1 protein stained by HDAC1 antibody (GTX100513) diluted at 1:500.


Red: phalloidin, a cytoskeleton marker, stained by phalloidin (invitrogen, A12380) diluted at 1:200.

Blue: Hoechst 33342 staining.



#### GTX100513 WB Image

HDAC1 antibody detects HDAC1 protein by western blot analysis. Various whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with HDAC1 antibody (GTX100513) diluted by 1:1000.




#### GTX100513 IHC-P Image

HDAC1 antibody detects HDAC1 protein at nucleus on mouse liver by immunohistochemical analysis.

Sample: Paraffin-embedded mouse liver.


HDAC1 antibody (GTX100513) dilution: 1:500.



#### GTX100513 IHC-P Image

HDAC1 antibody detects HDAC1 protein at nucleus on mouse colon by immunohistochemical analysis.

Sample: Paraffin-embedded mouse colon.  
HDAC1 antibody (GTX100513) dilution: 1:500.




#### GTX100513 IHC-P Image

HDAC1 antibody detects HDAC1 protein at cytoplasm and nucleus in rat brain by immunohistochemical analysis.


Sample: Paraffin-embedded rat brain.

HDAC1 antibody (GTX100513) diluted at 1:500.



#### GTX100513 WB Image

Various whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with HDAC1 antibody (GTX100513) diluted at 1:1000.




#### GTX100513 IHC-P Image

HDAC1 antibody detects HDAC1 protein at nucleus in human lung adenocarcinoma by immunohistochemical analysis.


Sample: Paraffin-embedded human lung adenocarcinoma.

HDAC1 antibody (GTX100513) diluted at 1:250.




**GTX100513 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded human testis, using HDAC1(GTX100513) antibody(10 µg/ml).




**GTX100513 IP Image**

HDAC1 antibody immunoprecipitates HDAC1 protein in IP experiments. IP Sample: 1000 µg 293T whole cell lysate/extract A. 40 µg 293T whole cell lysate/extract B. Control with 2.5 µg of preimmune rabbit IgG C. Immunoprecipitation of HDAC1 protein by 2.5 µg of HDAC1 antibody (GTX100513) 10% SDS-PAGE The immunoprecipitated HDAC1 protein was detected by HDAC1 antibody (GTX100513) diluted at 1:1000. EasyBlot anti-rabbit IgG (GTX221666-01) was used as a secondary reagent.




**GTX100513 ICC/IF Image**

Immunofluorescence analysis of paraformaldehyde-fixed A431, using HDAC1(GTX100513) antibody at 1:200 dilution.



**GTX100513 ChIP Image**

HDAC1 antibody immunoprecipitates HDAC1 protein-DNA in ChIP experiments. ChIP Sample: 293T whole cell lysate/extract A. 5 µg preimmune rabbit IgG B. 5 µg of HDAC1 antibody (GTX100513) The precipitated DNA was detected by PCR with primer set targeting to p21 promoter.




**GTX100513 WB Image**

Sample (30 ug of whole cell lysate)

A: NIH-3T3  
7.5% SDS PAGE

GTX100513 diluted at 1:1000



**GTX100513 WB Image**

HDAC1 antibody detects HDAC1 protein by Western blot analysis.

A. 30 µg Rat2 whole cell lysate/extract

10 % SDS-PAGE


HDAC1 antibody (GTX100513) dilution: 1:1000

### Application Reference

1. Lokireddy S (2015) *Proc Natl Acad Sci U S A* E7176-85
2. Lee KH (2014) *Sci Rep* 6394
3. Kumar S (2014) *Development* 2972-7
4. Chang JF (2014) *Toxicol Sci* 396-406
5. Yoon JH (2012) *Biochem Pharmacol* 747-57

### DataSheet - 5

<b>Catalog Number</b>	GTX213110-01	Package: 1 ml	<a href="#">Reference</a> ( 38 )
<b>Product Name</b>	Rabbit IgG antibody (HRP)		
<b>Synonyms</b>	antiRabbit IgG antibody, anti-Rabbit IgG antibody, Goat antiRabbit IgG HRP, Rabbit IgG secondary antibody, Goat anti-Rabbit IgG HRP, anti Rabbit IgG antibody, Goat anti Rabbit IgG HRP		
<b>Product Description</b>	HRP-conjugated Goat anti-Rabbit IgG polyclonal antibody		
<b>Background</b>	Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/ml in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.		
<b>Host</b>	Goat		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Target</b>	Rabbit IgG		
<b>Immunogen</b>	Highly purified whole rabbit IgG		
<b>Antigen Species</b>	Rabbit		
<b>Cross Reactivity Note</b>	Rabbit		
<b>Applications</b>	Dot, ELISA, IHC-P, WB		
<b>Application Note</b>		Suggested dilution	Reference
	Dot blot	Assay-dependent dilution	
	ELISA	Assay-dependent dilution	
	IHC (Formalin-fixed paraffin-embedded sections)	1:100-1:1000*	
	Western blot	Assay-dependent dilution	
	Not tested in other applications. *Optimal dilutions/concentrations should be determined by the researcher.		
<b>Conjugation</b>	HRP		
<b>Form Supplied</b>	Liquid		
<b>Purification</b>	Affinity purified with antigen		
<b>Concentration</b>	0.15 mg/ml (Please refer to the vial label for the specific concentration)		
<b>Storage Buffer</b>	0.05M Tris, 0.15M NaCl (pH7.4), 1%BSA. 0.025% ProCin 300 was added as a preservative.		
<b>Storage Instruction</b>	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.		
<b>Notes</b>	For <i>In vitro</i> laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.		




#### GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in mouse intestine by immunohistochemical analysis.

Sample: Paraffin-embedded mouse intestine.

WBP11 antibody (GTX118654) diluted at 1:500.

The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01)



#### GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in rat prostate by immunohistochemical analysis.

Sample: Paraffin-embedded rat prostate.

WBP11 antibody (GTX118654) diluted at 1:500.

The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01).

#### Application Reference

1. Lin CW (2016) *Acta Biomater*
2. Choi JY (2015) *Biochem Biophys Res Commun* 883-8
3. Pillay S (2016) *Nature*
4. Chuang YC (2014) *J Virol* 13759-68
5. Chen HY (2015) *Vet Parasitol* 281-91
6. Chun-Chang Yeh (2015) *BMC Anesthesiology* 92
7. Chen WH (2015) *Cancer Lett.* 65-74
8. Rumwald Leo G (2014) *SPIE Proceedings* 894405-1
9. Yang DF (2014) *J Formos Med Assoc.* 88-93
10. Huang JH (2015) *PLoS Pathog* e1004985
11. Lee WY (2015) *Oncotarget*
12. Eling N (2015) *Oncoscience* 517-32
13. Huang CY (2015) *Int J Mol Sci* 14171-80
14. Nils Eling (2015) *Oncoscience* 517-32
15. Kuan-Lin Kuo (2015) *Cancer Letters* 127–136
16. Heng-Hsiung Wu (2015) *Journal of Experimental Medicine* [Epub ahead of print]
17. *Int J Mol Med.* (2014)
18. Hsu TH (2014) *Cell Death Differ*
19. Chen HY (2014) *Vet Parasitol*
20. Zheng Y (2014) *PLoS One* e98552
21. Grossini E (2014) *J Endocrinol* 137-49
22. Wang HC (2014) *Eur J Med Chem* 312-34
23. Jagtap AD (2014) *Eur J Med Chem* 268-288
24. Wei MF (2014) *Autophagy* 1179-92
25. Tseng LC (2014) *PLoS One* e93394
26. Chen CM (2014) *Free Radic Biol Med*

27. Hsu CY (2014) *PLoS Pathog* e1003974
29. Kilgore JA (2013) *J Biol Chem* 19673-84
31. Chiang HC (2012) *Respiration* 319-26
33. Liu K (2012) *Gene* 225-30
35. Chang KH (2013) *Evid Based Complement Alternat Med* 471659
37. Lin YL (2012) *Brain Behav Immun* 459-68
28. Lee JG (2014) *Toxicol In Vitro* 562-570
30. Fu PK (2012) *Evid Based Complement Alternat Med* 837513
32. Xie H (2012) *PLoS One* e33087
34. Huang KH (2012) *PLoS One* e33615
36. Chen YH (2012) *PLoS One* e48335
38. Cheng AN (2013) *Cancer Lett* 218-25