

# Grid Screen™

## Sodium Chloride

HAMPTON  
RESEARCH

*Solutions for Crystal Growth*

### User Guide

HR2-219 (pg 1)

Grid Screen™ Sodium Chloride is a preformulated reagent kit designed to provide a rapid screening method for the crystallization of biological macromolecules. The screen is simple and practical for finding initial crystallization conditions as well as determining the solubility of a macromolecule in Sodium Chloride between pH 4.0 and 9.0.

### Sample Preparation

The macromolecular sample should be homogenous, as pure as is practically possible (>95%) and free of amorphous and particulate material. Remove amorphous material by centrifugation or micro-filtration prior to use.<sup>1,2,3</sup>

The recommended sample concentration is 5 to 25 mg/ml in water. Initially, the sample should be free of any unnecessary additives in order to observe the effect of the Grid Screen Sodium Chloride variables. Ideally, the initial screen should be performed with a sample which has been dialyzed against water although ligands, ions, reducing agents, or other additives may be present as required by the sample for solubility, stability, or activity.


### Performing the Screen

Since it is the most frequently reported method of crystallization, the following procedure describes the use of Grid Screen Sodium Chloride with the Hanging Drop Vapor Diffusion method. Grid Screen Sodium Chloride is also very compatible with the Sitting Drop, Sandwich Drop, MicroBatch, and Microdialysis methods. A complete description of the Hanging, Sitting, Sandwich Drop, Dialysis and other crystallization methods are available from the Hampton Research Crystal Growth 101 Library.

1. Prepare a VDX Plate (HR3-140) for Hanging Drop Vapor Diffusion by applying a thin bead of cover slide sealant to the upper edge of each of the 24 reservoirs. One may also use a Greased VDX Plate (HR3-170). Twenty-four reservoirs are to be prepared for a complete Grid Screen Sodium Chloride. See Figure 1.

**Figure 1**


Cross section of a reservoir in the VDX plate.



2. Using a clean pipet tip, pipet 1 ml of Grid Screen Sodium Chloride reagent A1 into reservoir A1. Discard the pipet tip, add a new pipet tip and pipet 1 ml of Grid Screen Sodium Chloride reagent A2 into reservoir A2. Repeat the procedure for the remaining 22 Grid Screen Sodium Chloride reagents using a clean pipet tip for each reagent so as to avoid reagent contamination and carry over.

3. Pipet 2  $\mu$ l of the sample to the center of a clean, siliconized 22 mm diameter circle or square cover slide. See Figure 2.

**Figure 2**




4. Pipet 2  $\mu$ l of Grid Screen Sodium Chloride reagent A1 from reservoir A1 into the sample droplet and mix by aspirating and dispensing the droplet several times, keeping the tip in the drop during mixing to avoid foaming. See Figure 2.

5. Working quickly to minimize evaporation, invert the cover slide and droplet over reservoir A1 and seal the cover slide onto the edge of the reservoir. See Figure 3.

**Figure 3**

Inverted Siliconized Coverslip placed over the reservoir.



6. Repeat operations 3 through 5 for the remaining 23 Grid Screen Sodium Chloride reagents.

7. If the quantity of sample permits, perform Grid Screen Sodium Chloride in duplicate and incubate one set of plates at 4°C and the second set at room temperature. Incubate and store the crystallization plates in a stable temperature environment free of vibration.

### Examine the Drop


Carefully examine the drops under a stereo microscope (10 to 100x magnification) immediately after setting up the screen. Record all observations and be particularly careful to scan the focal plane for small crystals. Observe the drops once each day for the first week, then once a week thereafter. Records should indicate whether the drop is clear, contains precipitate, and or crystals. It is helpful to describe the drop contents using descriptive terms. Adding magnitude is also helpful. Example: 4+ yellow/brown fine precipitate, 2+ small bipyramid crystals, clear drop, 3+ needle shaped crystals in 1+ white precipitate. One may also employ a standard numerical scoring scheme (Clear = 0, Precipitate = 1, Crystal = 10, etc). Figure 4 (on page 2) shows typical examples of what one might observe in a crystallization experiment.

### Interpreting Grid Screen Sodium Chloride


Clear drops indicate that either the relative supersaturation of the sample and reagent is too low or the drop has not yet completed equilibration. If

**Figure 4**


Typical observations in a crystallization experiment




Clear Drop




Skin/Precipitate




Precipitate




Precipitate/Phase




Quasi Crystals



Microcrystals




Needle Cluster



Plates



Rod Cluster



Single Crystal

the drop remains clear after 3 to 4 weeks consider repeating the Grid Screen Sodium Chloride condition and doubling the sample concentration. If more than 70% Grid Screen Sodium Chloride drops are clear consider doubling the sample concentration and repeating the entire screen.

Drops containing precipitate indicate that either the relative supersaturation of the sample and reagent is too high, the sample has denatured, or the sample is heterogeneous. To reduce the relative supersaturation, dilute the sample twofold and repeat the Grid Screen Sodium Chloride condition. If more than 70% Grid Screen Sodium Chloride drops contain precipitate and no crystals are present, consider diluting the sample concentration in half and repeating the entire screen. If sample denaturation is suspect, take measures to stabilize the sample (add reducing agent, ligands, glycerol, salt, or other stabilizing agents). If the sample is impure, aggregated, or heterogeneous take measures to pursue homogeneity. It is possible to obtain crystals from precipitate so do not discard nor ignore a drop containing precipitate. If possible, examine drops containing precipitate under polarizing optics to differentiate precipitate from microcrystalline material.

If the drop contains a macromolecular crystal the relative supersaturation of the sample and reagent is good. The next step is to optimize the preliminary conditions (pH, salt type, salt concentration, precipitant type, precipitant concentration, sample concentration, temperature, additives, and other crystallization variables) which produced the crystal in order to improve crystal size and quality.

Compare the observations between the 4°C and room temperature incubation to determine the effect of temperature on sample solubility. Different results in the same drops at different temperatures indicate that sample solubility is temperature dependent and that one should include temperature as a variable in subsequent screens and optimization experiments.

Retain and observe plates until the drops are dried out. Crystal growth can occur within 15 minutes or one year.

### Grid Screen Sodium Chloride Formulation

Grid Screen Sodium Chloride reagents are formulated using the highest purity chemicals, ultrapure water (18.2 Megohm-cm, 5 ppb TOC) and are sterile filtered using

0.22 micron filters into sterile containers (no preservatives added).

Grid Screen Sodium Chloride reagents are stable at room temperature and are best if used within 12 months of receipt. To enhance reagent stability it is strongly recommended that Grid Screen Sodium Chloride be stored at 4°C or -20°C. Avoid ultraviolet light to preserve reagent stability.

### References and Readings

1. Crystallization of nucleic acids and proteins, Edited by A. Ducruix and R. Giegé, The Practical Approach Series, Oxford Univ. Press, 1992.
2. Current approaches to macromolecular crystallization. McPherson, A. Eur. J. Biochem. 189, 1-23, 1990.
3. Protein and Nucleic Acid Crystallization. Methods, A Companion to Methods in Enzymology, Academic Press, Volume 1, Number 1, August 1990.
4. Advance in Protein Chemistry Volume 41. Pages 1-33 (Patricia C. Weber). Academic Press, 1991.
5. Current approaches to macromolecular crystallization., McPherson, A., Eur. J. Biochem. 189, 1-23, 1990.
6. Crystallization of Membrane Proteins. Edited by Hartmut Michel, 1990. CRC Press.

### Technical Support

Inquiries regarding Grid Screen Sodium Chloride reagent formulation, interpretation of screen results, optimization strategies and general inquiries regarding crystallization are welcome. Please e-mail, fax, or telephone your request to Hampton Research. Fax and e-mail Technical Support are available 24 hours a day. Telephone technical support is available 8:00 a.m. to 4:30 p.m. USA Pacific Standard Time.

Hampton Research  
34 Journey

Aliso Viejo, CA 92656-3317 U.S.A.

Tel: (949) 425-1321 • Fax: (949) 425-1611

Technical Support e-mail: tech@hrmail.com

Website: www.hamptonresearch.com

[ Sodium chloride (M) ]	pH					
	4	5	6	7	8	9
1.0	A1	A2	A3	A4	A5	A6
2.0	B1	B2	B3	B4	B5	B6
3.0	C1	C2	C3	C4	C5	C6
4.0	D1	D2	D3	D4	D5	D6

The pH indicated on each Grid Screen reagent is the ACTUAL pH of the reagent at 25.0 ° C. All pH adjustments have been made using Hydrochloric acid or Sodium hydroxide.

Tube #	Sodium chloride [ M ]	Tube #	Buffer
A1.	1.0	A1.	0.1 M Citric acid pH 4.0
B1.	2.0	B1.	0.1 M Citric acid pH 4.0
C1.	3.0	C1.	0.1 M Citric acid pH 4.0
D1.	4.0	D1.	0.1 M Citric acid pH 4.0
A2.	1.0	A2.	0.1 M Citric acid pH 5.0
B2.	2.0	B2.	0.1 M Citric acid pH 5.0
C2.	3.0	C2.	0.1 M Citric acid pH 5.0
D2.	4.0	D2.	0.1 M Citric acid pH 5.0
A3.	1.0	A3.	0.1 M MES monohydrate pH 6.0
B3.	2.0	B3.	0.1 M MES monohydrate pH 6.0
C3.	3.0	C3.	0.1 M MES monohydrate pH 6.0
D3.	4.0	D3.	0.1 M MES monohydrate pH 6.0
A4.	1.0	A4.	0.1 M HEPES pH 7.0
B4.	2.0	B4.	0.1 M HEPES pH 7.0
C4.	3.0	C4.	0.1 M HEPES pH 7.0
D4.	4.0	D4.	0.1 M HEPES pH 7.0
A5.	1.0	A5.	0.1 M Tris pH 8.0
B5.	2.0	B5.	0.1 M Tris pH 8.0
C5.	3.0	C5.	0.1 M Tris pH 8.0
D5.	4.0	D5.	0.1 M Tris pH 8.0
A6.	1.0	A6.	0.1 M BICINE pH 9.0
B6.	2.0	B6.	0.1 M BICINE pH 9.0
C6.	3.0	C6.	0.1 M BICINE pH 9.0
D6.	4.0	D6.	0.1 M BICINE pH 9.0

#### Chemical Analysis and Recommended Optimization Reagents

<b>HR2-637</b> - 5.0 M Sodium chloride, 200 milliliters	Mr 58.44	NaCl	CAS Number [7647-14-5]	EC No 231-598-3
<b>HR2-831</b> - 1.0 M Citric acid, 100 milliliters	Mr 192.13	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	CAS Number [77-92-9]	EC No 201-069-1
<b>HR2-587</b> - 0.5 M MES monohydrate, 100 milliliters	Mr 213.25	C <sub>6</sub> H <sub>13</sub> NO <sub>4</sub> S · H <sub>2</sub> O	CAS Number [145224-94-8]	EC No 224-632-3
<b>HR2-585</b> - 1.0 M HEPES, 100 milliliters	Mr 238.31	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub> S	CAS Number [7365-45-9]	EC No 230-907-9
<b>HR2-589</b> - 1.0 M Tris, 100 milliliters	Mr 121.14	C <sub>4</sub> H <sub>11</sub> NO <sub>3</sub>	CAS Number [77-86-1]	EC No 201-064-4
<b>HR2-509</b> - 1.0 M BICINE, 100 milliliters	Mr 163.17	C <sub>6</sub> H <sub>13</sub> NO <sub>4</sub>	CAS Number [150-25-4]	EC No 205-755-1

Sample: \_\_\_\_\_ Sample Concentration: \_\_\_\_\_  
 Sample Buffer: \_\_\_\_\_ Date: \_\_\_\_\_  
 Reservoir Volume: \_\_\_\_\_ Temperature: \_\_\_\_\_  
 Drop Volume: Total \_\_\_\_\_  $\mu$ l Sample \_\_\_\_\_  $\mu$ l Reservoir \_\_\_\_\_  $\mu$ l Additive \_\_\_\_\_  $\mu$ l

- |   |  |
|---|--|
| 1 Clear Drop                                | 5 Posettes or Spherulites              |
| 2 Phase Separation                          | 6 Needles (1D Growth)                  |
| 3 Regular Granular Precipitate              | 7 Plates (2D Growth)                   |
| 4 Birefringent Precipitate or Microcrystals | 8 Single Crystals (3D Growth < 0.2 mm) |
|   | 9 Single Crystals (3D Growth > 0.2 mm) |

Grid Screen™ NaCl - HR2-219 Scoring Sheet	Date:	Date:	Date:	Date:
A1. 0.1 M Citric acid pH 4.0, 1.0 M Sodium chloride				
A2. 0.1 M Citric acid pH 5.0, 1.0 M Sodium chloride				
A3. 0.1 M MES monohydrate pH 6.0, 1.0 M Sodium chloride				
A4. 0.1 M HEPES pH 7.0, 1.0 M Sodium chloride				
A5. 0.1 M Tris pH 8.0, 1.0 M Sodium chloride				
A6. 0.1 M BICINE pH 9.0, 1.0 M Sodium chloride				
B1. 0.1 M Citric acid pH 4.0, 2.0 M Sodium chloride				
B2. 0.1 M Citric acid pH 5.0, 2.0 M Sodium chloride				
B3. 0.1 M MES monohydrate pH 6.0, 2.0 M Sodium chloride				
B4. 0.1 M HEPES pH 7.0, 2.0 M Sodium chloride				
B5. 0.1 M Tris pH 8.0, 2.0 M Sodium chloride				
B6. 0.1 M BICINE pH 9.0, 2.0 M Sodium chloride				
C1. 0.1 M Citric acid pH 4.0, 3.0 M Sodium chloride				
C2. 0.1 M Citric acid pH 5.0, 3.0 M Sodium chloride				
C3. 0.1 M MES monohydrate pH 6.0, 3.0 M Sodium chloride				
C4. 0.1 M HEPES pH 7.0, 3.0 M Sodium chloride				
C5. 0.1 M Tris pH 8.0, 3.0 M Sodium chloride				
C6. 0.1 M BICINE pH 9.0, 3.0 M Sodium chloride				
D1. 0.1 M Citric acid pH 4.0, 4.0 M Sodium chloride				
D2. 0.1 M Citric acid pH 5.0, 4.0 M Sodium chloride				
D3. 0.1 M MES monohydrate pH 6.0, 4.0 M Sodium chloride				
D4. 0.1 M HEPES pH 7.0, 4.0 M Sodium chloride				
D5. 0.1 M Tris pH 8.0, 4.0 M Sodium chloride				
D6. 0.1 M BICINE pH 9.0, 4.0 M Sodium chloride				