*ImmunoReagents,Inc.

840 Main Campus Drive Suite 3520 Raleigh, NC 27606

Product Specifications

Description:Goat anti-Human IgA (α chain) - Affinity Pure, FITC ConjugatePart Number:GtxHu-001-DFTTCConcentration:1.50 mg/ml (E 1% at 280 nm = 13.0)Amount:1.0 mgConjugate:Fluorescein-5-isothiocyanate (FTTC) Amax = 494 nm; Emax = 518 nmForm:Clear, fluorescein yellow liquidPurification:Affinity purified using solid phase Human IgAPurify:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2.8 °CSheef Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, his antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:Fils chains on all human immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:- 1:20 - 1:2,000 for most applicationsPischaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Product:	Goat anti-Human IgA (α chain) - Affinity Pure
Concentration:1.50 mg/ml (£ 1% at 280 nm = 13.0)Amount:1.0 mgConjugate:Fluorescein-5-isothiocyanate (FITC) Amax = 494 nm; Emax = 518 nmForm:Clear, fluorescent yellow liquidPurify:Affinity purified using solid phase Human IgAPurity:Affinity purified anibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.discard.assed on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:- 1:20-1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a licenser	Description:	Goat anti-Human IgA (α chain) - Affinity Pure, FITC Conjugate
Amount:1.0 mgConjugate:Fluorescein-5-isothiocyanate (FITC) Amax = 494 nm; Emax = 518 nmForm:Clear, fluorescent yellow liquidPurification:Affinity purified using solid phase Human IgAPurify:Affinity purified using solid phase Human IgAPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Proteasc/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):Disclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Part Number:	GtxHu-001-DFITC
Conjugate:Fluorescein-5-isothiocyanate (FTTC) Amax = 494 nm; Emax = 518 nmForm:Clear, fluorescent yellow liquidPurification:Affinity purified using solid phase Human IgAPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Concentration:	1.50 mg/ml (E 1% at 280 nm = 13.0)
Form:Clear, fluorescent yellow liquidPurification:Affinity purified using solid phase Human IgAPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Amount:	1.0 mg
Purification:Affinity purified using solid phase Human IgAPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • light chains on all human immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Conjugate:	Fluorescein-5-isothiocyanate (FITC) Amax = 494 nm; Emax = 518 nm
Purity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Form:	Clear, fluorescent yellow liquid
Host:GoatImmunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • ight chains on all human immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):Disclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products ar not recommendations to use our products in violation of any patent or as a license	Purification:	Affinity purified using solid phase Human IgA
Immunogen:Purified Human IgA, (α chain)Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • light chains on all human immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):Disclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Purity:	Affinity purified antibody is $> 95\%$ based on SDS-PAGE
Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglob	Host:	Goat
Protease/IgG freePreservative:0.05% (w/v) Sodium AzideStorage:2-8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Immunogen:	Purified Human IgA, (α chain)
Preservative:0.05% (w/v) Sodium AzideStorage:2.8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):Disclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA,
Preservative:0.05% (w/v) Sodium AzideStorage:2.8 °CShelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):Disclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		Protease/IgG free
Shelf Life:1 year from date of receipt. Prepare working dilution prior to use and then discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): • 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Preservative:	
discard.Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): • 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Storage:	2-8 °C
Specificity:Based on IEP, this antibody reacts with: • heavy (α) chains on human IgACross Reactivity:Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): • 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Shelf Life:	1 year from date of receipt. Prepare working dilution prior to use and then
 heavy (α) chains on human IgA Cross Reactivity: Based on IEP, no reactivity is observed to: non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins light chains on all human immunoglobulins Country of Origin: Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. Applications: This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license 		discard.
Cross Reactivity:Based on IEP, no reactivity is observed to:• non-immunoglobulin human serum immunoglobulins• light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Specificity:	Based on IEP, this antibody reacts with:
Cross Reactivity:Based on IEP, no reactivity is observed to:• non-immunoglobulin human serum immunoglobulins• light chains on all human immunoglobulinsCountry of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		• heavy (a) sheins on human Is A
 non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins light chains on all human immunoglobulins Country of Origin: Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. Applications: This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license 		• neavy (α) chains on numan 1gA
 light chains on all human immunoglobulins Country of Origin: Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. Applications: This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license 	Cross Reactivity:	Based on IEP, no reactivity is observed to:
Country of Origin:Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		• non-immunoglobulin human serum immunoglobulins
Applications:a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		 light chains on all human immunoglobulins
Applications:a registered veterinarian.Applications:This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s):• 1:20 - 1:2,000 for most applicationsDisclaimer:For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Country of Origin:	Goat serum was obtained from healthy animals of US origin and under the care of
 working dilution should be determined by the investigator. Suggested starting dilution(s): 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license 		
dilution(s): • 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license	Applications:	This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal
 1:20 - 1:2,000 for most applications Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license 		working dilution should be determined by the investigator. Suggested starting
Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		dilution(s):
Disclaimer: For <i>in vitro</i> Laboratory Use Only. Not intended for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		• $1.20 - 1.2000$ for most applications
Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license		
not recommendations to use our products in violation of any patent or as a license	Disclaimer:	For in vitro Laboratory Use Only. Not intended for diagnostic or therapeutic use.
		Not for human or animal consumption. Suggested applications of our products are
		not recommendations to use our products in violation of any patent or as a license
under any patent of ImmunoReagents, Inc.		under any patent of ImmunoReagents, Inc.
Phone: 919-831-2240 Fax: 919-831-2240 email: info@immunoreagents.com	Phone: 919-831-2240	Fax: 919-831-2240 email: info@immunoreagents.com

Product may not be resold or modified for resale without prior written approval of ImmunoReagents, Inc.