



KBI-60004 Tissue Digestion Kit II

For heavily cross-linked paraffin-embedded tissues

Instructions for use for KBI-60004 Tissue Digestion Kit II in combination with Kreatech™ FISH probes.

Fluorescent *in situ* hybridization (FISH) identifies or labels target genomic sequences so that their location can be studied. DNA sequences from appropriate chromosome specific probes are first labeled with reporter molecules. The labeled DNA probe is then hybridized to the metaphase chromosomes or interphase nuclei on a slide. After washing, the specimen is screened for the reporter molecules by fluorescence microscopy.

This pretreatment kit is specifically developed to obtain optimal results on heavily cross-linked or difficult paraffin embedded tissues (e.g. longer fixation or prolonged storage).

Note: Conventional paraffin embedded tissues

The use of the [KBI-60007 Tissue Digestion Kit I](#) is recommended for use with conventional samples.

For metaphase and interphase cells or other cytological samples it is recommended to use pretreatment kit KBI-60005 or KBI-60006.

For more info consult our website: www.leicabiosystems.com

Pretreatment:

Mount 4 - 6 µm formalin-fixed paraffin-embedded tissue sections on positively charged slides (e.g. aminoalkylsilane)

1. Bake mounted slides for 2 hours at 80 °C or 16 hours at 56 °C.
2. De-paraffinize warm slides by soaking in xylene or xylene substitute for two times 10 minutes (min).
3. Re-hydrate by soaking in 100%, 85% and 70% ethanol for 3 min each.
4. Wash with dH₂O for 3 min at room temperature (RT).
5. Pretreat with 0.2 M HCl (LK-099B) for 20 min, and then wash in dH₂O for 3 min at RT.
6. Place slides in Pretreatment Solution B (8% sodium thiocyanate) (LK-100B) at 80 °C for 30 min.
7. Rinse in 2 x SSC (LK-104B) for 3 min at RT.
8. Cover paraffin section with approximately 200 µl **Pepsin Solution (LK-101B) and incubate at RT for 5 - 50 min** (Time depending on tissue fixation and tissue type. E.g. most breast cancer tissue needs 5 - 15 min digestion; colon tissue 20 - 30 min).
9. Wash in dH₂O for 1 min
10. Wash in 2 x SSC (LK-104B) for 5 min at RT.
11. Dehydrate slides by soaking 70%, 85%, and 100% ethanol for 1 min each time. Air-dry. Proceed with **Probe preparation**.

Note: Check protein digestion and pretreatment by applying 15 µl DAPI counterstain and evaluate slides using a fluorescence microscope equipped with a DAPI filter. Remove cover slip and soak tissue in 2 x SSC for 2 min and prolong protein digestion if sample is not sufficiently digested. Use a fresh sample and reduce protein digestion time if the sample is over-digested.

Probe preparation:

Kreatech™ FISH probes for paraffin embedded tissue are supplied Ready to Use (RtU) unless specified otherwise in the product documentation. Consult label on vial and specific probe pack insert for dilution specifics.

Co-denaturation:

Apply 10 µl of probe per 22 x 22 mm field. Cover with glass cover slip and seal with Fixogum or rubber cement. Denature sample and probe on a *Thermobrite* (TS-01-02) at 80 ±1 °C for 5 min. Continue with **hybridization**.

Hybridization:

Incubate overnight at 37 ±1 °C in a *Thermobrite* (TS-01/02) or in a humidified chamber.

Post-Hybridization Wash:

1. Pre-warm Wash Buffer I (0.4 x SSC / 0.3% Igepal) (LK-102) to 72 °C
2. Remove rubber cement.
3. Place up to 14 slides in 200 ml of Wash Buffer II (2 x SSC / 0.1% Igepal) (LK-103), incubate for 2 min. at RT. Slide off coverclips. Re-use only once for a total of 28 slides.
4. Place up to 14 slides in 200 ml of pre-warmed Wash Buffer I (0.4 x SSC / 0.3% Igepal) (LK-102), incubate for 2 min at 72 °C (+- 1 °C) without agitation. Re-use only once for a total of 28 slides.
5. Place up to 14 slides in 200 ml of fresh Wash Buffer II (2 x SSC / 0.1% Igepal) (LK-103), incubate for 1 min. at RT without agitation. Re-use only once for a total of 28 slides.
6. Dehydrate in fresh 70%, 85% and 100% ethanol, incubate for 1 min each at RT. Air dry at RT and proceed to **counterstaining**.

Counterstaining:

Apply 15 µl DAPI counterstain (DAPI/Antifade 1 µg/ml) (LK-096B) and apply glass cover slip. Proceed with microscopy.

Procedural recommendations:

Temperature and buffer concentration (stringency) of hybridization and washing are important, as lower stringency can result in non-specific binding of the probe to other sequences, and higher stringency can result in a lack of signal. Incomplete denaturation of target DNA can result in lack of signal.

Recommendations for Fluorescence Microscopy:

For optimal visualization use a well maintained and regularly calibrated microscope equipped with a 100 W mercury lamp and a 63x or 100x fluorescent objective. Triple band-pass filters (DAPI/FITC/Texas Red or DAPI/FITC/Rhodamine) are used to view multiple colors, single band-pass filters are used for individual color visualization.

Suitable excitation and emission range for Kreatech™ fluorophores:

PlatinumBright™415 Ex 415 ±20 nm Em 475 ±30 nm



PlatinumBright™495 Ex 495 ±20 nm Em 525 ±30 nm

PlatinumBright™550 Ex 546 ±12 nm Em 580 ±30 nm

Warnings and Precautions:

- For *in vitro* diagnostic use.** For professional use only. In case of emergencies check SDS sheets for safety information.
- DNA probes and hybridization buffers contain formamide which is a teratogen; do not inhale or allow skin contact. Wear gloves and a lab coat when handling DNA probes and DAPI counterstain.
- All materials should be disposed of according to your institution's guidelines for hospital waste disposal.

Labelling According Regulation (EC) No 1272/2008

Code #	Description	Signal Word	Pictogram	Hazard Statements
LK-096B	DAPI/Antifade 1 µg/ml (DAPI Counterstain)	Warning		H315, H319, H335
LK-097B	Antifade (Counterstain diluent)	Warning		H315, H319, H335
LK-099B	0.2 M HCl	N.A.	N.A.	N.A.
LK-100B	8% sodium thiocyanate (Pretreatment Solution B)	N.A.	N.A.	N.A.
LK-101B	Pepsin Solution	N.A.	N.A.	N.A.
LK-102B	0.4 x SSC / 0.3% Igepal (Wash Buffer I)	N.A.	N.A.	N.A.
LK-103B	2 x SSC / 0.1% Igepal (Wash Buffer II)	N.A.	N.A.	N.A.
LK-104B	2 x SSC	N.A.	N.A.	N.A.

Material provided:

Code #	Description	Volume
LK-096B	DAPI/Antifade 1 µg/ml (DAPI Counterstain)	0.5 ml
LK-097B	Antifade (Counterstain diluent)	0.5 ml
LK-099B	0.2 M HCl	250 ml
LK-100B	8% sodium thiocyanate (Pretreatment Solution B)	250 ml
LK-101B	Pepsin Solution	5 ml
LK-102B	0.4 x SSC / 0.3% Igepal (Wash Buffer I)	250 ml
LK-103B	2 x SSC / 0.1% Igepal (Wash Buffer II)	2 x 250 ml
LK-104B	2 x SSC	2 x 250 ml

(*Please note that LK-102 and LK-103 products containing the same number but different letters are identical except for the volume.

A= 100 ml, B=250 ml)

Material required, but not supplied:

- Xylene
- Ethanol 100%, 85% and 70%
- Fixogum (LK-071A) or rubber cement
- Hot plate with accurate temperature control up to 80 °C or ThermoBrite™ (TS01/TS02)
- Incubator with accurate temperature range of 37 - 56 °C
- Water bath with accurate temperature range of 37 - 80 °C
- Plastic or glass coplin jars
- Variable micropipettes (1 µl - 200 µl)
- Fluorescence microscope equipped with suitable filters (see recommendations for Fluorescence Microscopy).



**Kreatech
Biotechnology B.V.**
Vlierweg 20
1032 LG Amsterdam
The Netherlands

Patents & Tradenames:

Kreatech™ is a trade name of Kreatech Biotechnology B.V. REPEAT FREE™ is a trademark of Veridex LLC. All third party trademarks are the property of their respective owners. Leica Biosystems reserves the right to amend specifications without notice.

All Rights Reserved. © Leica Biosystems Richmond, Inc. 2013. The Kreatech™ products or the use of the Kreatech™ products is subject to intellectual property rights. FISH and CISH probes are produced under an exclusive license from Veridex LLC for its REPEAT-FREE™ technology. FISH probes are labeled with the Kreatech's proprietary Universal Linkage system (ULS™). The following fluorophores (-415; -547; -590; -647) are subject to patents and know-how, owned or controlled, and manufactured by DYOMICS GmbH. The ULS™ technology and products are covered by one or more of the following US patents and corresponding patent claims outside the US owned by Kreatech: US 5,580, 990; 5,714,327; 5,985,566; 6,133,038; US RE 40,557E, 6,797,818; 7,217,813.