


TACR2 / Nk2 / NK2R Rabbit anti-Human Polyclonal (Extracellular Domain) Antibody - LS-A1290 - LSBio	
CatalogID:	LS-A1290
Target:	tachykinin receptor 2 (TACR2)
Synonyms:	TACR2 Antibody, Neurokinin 2 receptor Antibody, NK-2 receptor Antibody, Neurokinin A receptor Antibody, NK2 Antibody, NK2R Antibody, Tachykinin receptor 2 Antibody, Substance-K receptor Antibody, NK-2R Antibody, SK receptor Antibody, SKR Antibody, TAC2R Antibody, NKNAR Antibody, Substance K receptor Antibody
Family / Subfamily:	GPCR / Tachykinin
Host	TACR2 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	TACR2 / Nk2 / NK2R antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	TACR2 / Nk2 / NK2R antibody was raised against synthetic 16 amino acid peptide from 2nd extracellular domain of human Neurokinin A Receptor. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset (100%); Bovine, Rabbit (94%); Horse, Pig (88%); Mouse, Rat, Bat, Hamster, Elephant, Guinea pig (81%).
Specificity:	Human Neurokinin A Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Epitope:	Extracellular Domain
Reactivity:	Human, Gorilla, Gibbon, Monkey
Predicted Reactivity:	Bovine, Rabbit
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Uses:	IHC - Paraffin (5 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-Neurokinin A Receptor antibody LS-A1290 IHC of human skeletal muscle.
Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences