



CDD116 Pabbi	t anti-Human Polyclonal (Internal) Antibody - LS-A1518 - LSBio
CatalogID:	LS-A1518
Target:	G protein-coupled receptor 116 (GPR116)
Synonyms:	GPR116 Antibody, Ig-Hepta homolog Antibody, Ig-hepta Antibody, G protein-coupled receptor 116 Antibody, KIAA0758 Antibody, KPG_001 Antibody
Family / Subfamily:	GPCR / Orphan-B
Host	GPR116 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	GPR116 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	GPR116 antibody was raised against synthetic 20 amino acid peptide from internal region of human GPR116. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon (100%); Marmoset (95%); Monkey (90%).
Specificity:	Human GPR116. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Epitope:	Internal
Reactivity:	Human, Gorilla, Gibbon
Predicted Reactivity:	Monkey
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Usage Summary:	Immunohistochemistry: LS-A1518 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after proteinase K antigen retrieval. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS -A1518 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 μg/ml), ELISA (Optimal dilution to be determined by the researcher)
Size:	50 μg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-GPR116 antibody LS-A1518 IHC of human heart. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Immunohistochemistry Image:



Anti-GPR116 antibody LS-A1518 IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences