


TACR3 / NK3R Rabbit anti-Human Polyclonal (C-Terminus) Antibody - LS-A4147 - LSBio	
CatalogID:	LS-A4147
Target:	tachykinin receptor 3 (TACR3)
Synonyms:	TACR3 Antibody, HH11 Antibody, Nmk-r Antibody, NK3R Antibody, NKR Antibody, Tachykinin receptor 3 Antibody, Neurokinin B Antibody, Neurokinin B receptor Antibody, NK-3 receptor Antibody, NK-3R Antibody, TAC3R Antibody, Neurokinin 3 receptor Antibody, Neurokinin beta receptor Antibody, Neuromedin-K receptor Antibody, TAC3RL Antibody
Family / Subfamily:	GPCR / Tachykinin
Host	TACR3 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	TACR3 / NK3R antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	TACR3 / NK3R antibody was raised against synthetic 17 amino acid peptide from C-terminal cytoplasmic domain of human TACR3. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey (100%); Gibbon, Marmoset (94%); Mouse, Rat, Hamster, Panda, Bovine, Dog, Rabbit (88%); Elephant, Bat (82%).
Specificity:	Human TACR3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Epitope:	C-Terminus
Reactivity:	Human, Gorilla
Predicted Reactivity:	Gibbon, Monkey
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Uses:	IHC - Paraffin (20 μg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 μg
Concentration:	1 mg/ml

## Immunohistochemistry Image:



Anti-TACR3 antibody LS-A4147 IHC of human brain, hippocampus. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences