


FZD7 / Frizzled 7 Rabbit anti-Human Polyclonal (N-Terminus) Antibody - LS-A4899 - LSBio	
CatalogID:	LS-A4899
Target:	frizzled class receptor 7 (FZD7)
Synonyms:	FZD7 Antibody, Frizzled family receptor 7 Antibody, Frizzled homolog 7 Antibody, Frizzled 7 Antibody, Frizzled-7 Antibody, FzE3 Antibody, Fz-7 Antibody, HFz7 Antibody
Family / Subfamily:	GPCR / Frizzled
Host	FZD7 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	FZD7 / Frizzled 7 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	FZD7 / Frizzled 7 antibody was raised against synthetic 20 amino acid peptide from N-terminal extracellular domain of human FZD7 / Frizzled 7. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bovine, Bat, Rabbit, Pig, Opossum (100%); Xenopus (90%); Turkey, Chicken (85%).
Specificity:	Human FZD7 / Frizzled 7. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FZD1 (60%), FZD2 (60%).
Epitope:	N-Terminus
Reactivity:	Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Bat, Bovine, Dog, Hamster, Pig, Rabbit
Predicted Reactivity:	Xenopus
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Uses:	IHC - Paraffin (4 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-FZD7 / Frizzled 7 antibody LS-A4899 IHC of human heart, cardiac myocytes.
Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences