


FZD9 / Frizzled 9 Rabbit anti-Human Polyclonal (Extracellular Domain) Antibody - LS-A5081 - LSBio	
<b>CatalogID:</b>	LS-A5081
<b>Target:</b>	frizzled class receptor 9 (FZD9)
<b>Synonyms:</b>	FZD9 Antibody, CD349 Antibody, Frizzled homolog 9 Antibody, Fz-9 Antibody, Frizzled family receptor 9 Antibody, Frizzled-9 Antibody, Frizzled 9 Antibody, Fz9 Antibody, HFz9 Antibody, CD349 antigen Antibody, Frizzled homolog fzd3 Antibody, FzE6 Antibody
<b>Family / Subfamily:</b>	GPCR / Frizzled
<b>Host</b>	FZD9 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	FZD9 / Frizzled 9 antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	FZD9 / Frizzled 9 antibody was raised against synthetic 14 amino acid peptide from 1st extracellular domain of human FZD9 / Frizzled 9. Percent identity with other species by BLAST analysis: Human, Gibbon, Mouse, Rat, Hamster, Panda, Dog, Bovine, Bat, Rabbit, Opossum (100%); Platypus (86%).
<b>Specificity:</b>	Human FZD9 / Frizzled 9. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
<b>Epitope:</b>	Extracellular Domain
<b>Reactivity:</b>	Human, Gibbon, Mouse, Rat, Bat, Bovine, Dog, Hamster, Rabbit
<b>Purification:</b>	Immunoaffinity purified
<b>Presentation:</b>	PBS, 0.1% sodium azide.
<b>Recommended Storage:</b>	Long term: -70°C; Short term: +4°C
<b>Uses:</b>	IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**



Anti-FZD9 / Frizzled 9 antibody LS-A5081 IHC of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences