


NR0B1 / DAX1 Rabbit anti-Human Polyclonal (N-Terminus) Antibody - LS-A5703 - LSBio	
<b>CatalogID:</b>	LS-A5703
<b>Target:</b>	nuclear receptor subfamily 0, group B, member 1 (NR0B1)
<b>Synonyms:</b>	NR0B1 Antibody, AHX Antibody, AHC Antibody, AHCH Antibody, DAX1 Antibody, DSS Antibody, DAX 1 Antibody, DAX-1 Antibody, Adrenal hypoplasia congenita Antibody, Dosage-sensitive sex reversal Antibody, GTD Antibody, HHG Antibody, Nuclear hormone receptor Antibody, NROB1 Antibody, Nuclear receptor DAX-1 Antibody, SRXY2 Antibody, Nuclear receptor DAX1 Antibody, Dax-1 receptor Antibody
<b>Family / Subfamily:</b>	NHR / NR0 Knirps-like
<b>Host</b>	NR0B1 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	NR0B1 / DAX1 antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	NR0B1 / DAX1 antibody was raised against synthetic 17 amino acid peptide from N-terminus of human DAX1. Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Marmoset (100%); Monkey (94%).
<b>Specificity:</b>	Human DAX1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
<b>Epitope:</b>	N-Terminus
<b>Reactivity:</b>	Human, Chimpanzee, Gorilla, Orangutan, Gibbon
<b>Predicted Reactivity:</b>	Monkey
<b>Purification:</b>	Immunoaffinity purified
<b>Presentation:</b>	PBS, 0.1% sodium azide.
<b>Recommended Storage:</b>	Long term: -70°C; Short term: +4°C
<b>Usage Summary:</b>	Immunohistochemistry: LS-A5703 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-A5703 was determined to be 10 ug/ml.
<b>Uses:</b>	IHC - Paraffin (10 µg/ml), ELISA (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	0.7 mg/ml

**Immunohistochemistry Image:**



Anti-DAX1 antibody LS-A5703 IHC of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences