


CatalogID:	LS-A822
Target:	cholecystokinin A receptor (CCKAR)
Synonyms:	CCKAR Antibody, CCK-A receptor Antibody, CCK-AR Antibody, CCKRA Antibody, CCK-A Antibody, CCKA receptor Antibody, Cholecystokinin A receptor Antibody, Cck1r Antibody, CCK1-R Antibody, Cholecystokinin-1 receptor Antibody
Family / Subfamily:	GPCR / Cholecystokinin
Host	CCKAR antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	CCKAR / CCK1R antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	CCKAR / CCK1R antibody was raised against synthetic 16 amino acid peptide from internal region of human CCKAR. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Dog, Panda (100%) Bovine, Elephant, Horse, Rabbit (94%); Bat, Opossum, Guinea pig (88%); Mouse, Rat, Hamster (81%).
Specificity:	Human CCKAR. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Epitope:	Internal
Reactivity:	Human, Gorilla, Gibbon, Monkey, Dog
Predicted Reactivity:	Bovine, Horse, Rabbit
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Usage Summary:	Immunohistochemistry with formalin-fixed paraffin-embedded tissues requires pretreatment with Proteinase K.
Uses:	IHC - Paraffin (5 - 10 μ g/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-CCKAR antibody LS-A822 IHC of human prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Immunohistochemistry Image:

Anti-CCKAR antibody	For the set of th
Requested From:	Japan
	atory Reagent For In Vitro Research Use Only
Not for resale wit	hout prior written consent from LifeSpan BioSciences, Inc.
	Created on 9/23/2014
	© 2014 LifeSpan BioSciences