


| wingless-type MMTV integration site family, member 4 (WNT4) whomps: WNT4 Antibody, WNT-4 Antibody, Protein Wnt-4 Antibody, SERKAL Antibody will / Subfamily: WNT / not assigned-WNT WNT4 antibody was produced in Rabbit Polyclonal WNT4 antibody was raised against Human MIGHT April Apri | | |
|--|--|---|
| wingless-type MMTV integration site family, member 4 (WNT4) whomps: WNT4 Antibody, WNT-4 Antibody, Protein Wnt-4 Antibody, SERKAL Antibody will / Subfamily: WNT / not assigned-WNT WNT4 antibody was produced in Rabbit Polyclonal WNT4 antibody was raised against Human MIGHT April Apri | WNT4 Rabbit anti-Human Polyclonal (Internal) Antibody - LS-A9622 - LSBio | |
| wnonyms: wno | CatalogID: | LS-A9622 |
| wntily / Subfamily: wntily / Subfamily: wntile antibody was produced in Rabbit wnunogen Species: wntile antibody was raised against Human wnunogen Species: wntile antibody was raised against Human synthetic peptide wntile antibody was raised against synthetic 14 amino acid peptide from internative region of human wntile percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). wecificity: Human wntile BLAST analysis of the peptide immunogen showed no homology with other human proteins, except wntile immunogen showed no homology with other human proteins, except wntile (17%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified wesentation: pBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Target: | wingless-type MMTV integration site family, member 4 (WNT4) |
| WNT4 antibody was produced in Rabbit Polyclonal WNT4 antibody was raised against Human MNT4 antibody was raised against Human Synthetic peptide WNT4 antibody was raised against synthetic 14 amino acid peptide from internategion of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Muman WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Infication: Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Synonyms: | WNT4 Antibody, WNT-4 Antibody, Protein Wnt-4 Antibody, SERKAL Antibody |
| Polyclonal Munogen Species: WNT4 antibody was raised against Human Synthetic peptide WNT4 antibody was raised against synthetic 14 amino acid peptide from internar region of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig urification: Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Family / Subfamily: | WNT / not assigned-WNT |
| wnunogen Species: WNT4 antibody was raised against Human Synthetic peptide WNT4 antibody was raised against synthetic 14 amino acid peptide from internaregion of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Host | WNT4 antibody was produced in Rabbit |
| Synthetic peptide WNT4 antibody was raised against synthetic 14 amino acid peptide from internate region of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Irrification: Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Clonality: | Polyclonal |
| WNT4 antibody was raised against synthetic 14 amino acid peptide from interna region of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Decificity: Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Immunogen Species: | WNT4 antibody was raised against Human |
| region of human WNT4. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Pig (100%); Opossum (86%). Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Antigen Type: | Synthetic peptide |
| with other human proteins, except WNT5A (71%). Internal Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified PBS, 0.1% sodium azide. PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Immunogen: | Human, Gorilla, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, |
| Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Specificity: | Human WNT4. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except WNT5A (71%). |
| Immunoaffinity purified PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Epitope: | Internal |
| PBS, 0.1% sodium azide. Long term: -70°C; Short term: +4°C IHC - Paraffin (10 µg/ml) (Optimal dilution to be determined by the researcher) | Reactivity: | Human, Gorilla, Monkey, Mouse, Rat, Bovine, Dog, Hamster, Horse, Pig |
| Long term: -70°C; Short term: +4°C IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Purification: | Immunoaffinity purified |
| Ses: IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) | Presentation: | PBS, 0.1% sodium azide. |
| | Recommended Storage: | Long term: -70°C; Short term: +4°C |
| ze : 50 μg | Uses: | IHC - Paraffin (10 μg/ml) (Optimal dilution to be determined by the researcher) |
| | Size: | 50 µg |
| oncentration: 1 mg/ml | Concentration: | 1 mg/ml |

Immunohistochemistry Image:



Anti-WNT4 antibody LS-A9622 IHC of human thyroid. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences