



| • • • • | |
|----------------------|---|
| CatalogID: | LS-B10821 |
| Validation: | This antibody replaces catalog number LS-C185735. It has been validated for use in the following assays: IHC-P. |
| Target: | KRR1, small subunit (SSU) processome component, homolog (yeast) |
| Synonyms: | KRR1 Antibody, HIV-1 Rev binding protein 2 Antibody, HIV-1 Rev-binding protein 2 Antibody, Rev-interacting protein 1 Antibody, Rev interacting protein Antibody, RIP 1 Antibody, HRB2 Antibody |
| Host | KRR1 antibody was produced in Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Immunogen Species: | RIP-1 / KRR1 antibody was raised against Human |
| Antigen Type: | Recombinant protein |
| Immunogen: | RIP-1 / KRR1 antibody was raised against recombinant fragment corresponding to a region within amino acids 47 and 344 of KRR1 (SwissProt Q13601). Percent identity by BLAST analysis: Human (100%); Mouse, Bovine (97%); Xenopus (87%); Chicken (85%). |
| Specificity: | Human RIP-1 / KRR1 |
| Epitope: | aa47-344 |
| Reactivity: | Human |
| Purification: | Immunoaffinity purified |
| Presentation: | 0.1 M Tris-glycine, pH 7.0, 10% glycerol, 0.01% Thimerosal |
| Recommended Storage: | Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Usage Summary: | IHC-paraffin: Suggested antigen retrieval using heat mediated 10 mM Citrate buffer (pH 6.0) or Tris-EDTA buffer (pH 8.0). |
| Uses: | IHC - Paraffin (7.5 μg/ml), ICC (1:100 - 1:1000), Western blot (1:500 - 1:3000) (Optimal dilution to be determined by the researcher) |
| Size: | 50 µl |
| Concentration: | 1 mg/ml |

Immunohistochemistry Image:



Human Kidney: Formalin-Fixed, Paraffin-Embedded (FFPE)

Immunocytochemistry Image:



Immunofluorescence of paraformaldehyde-fixed HeLa using KRR1 antibody at 1:500 dilution.

