



| ASNS Rabbit anti-Human Polyclonal Antibody - LS-B10899 - LSBio | |
|--|---|
| CatalogID: | LS-B10899 |
| Validation: | This antibody replaces catalog number LS-C155365. It has been validated for use in the following assays: IHC-P. |
| Target: | asparagine synthetase (glutamine-hydrolyzing) (ASNS) |
| Synonyms: | ASNS Antibody, Asparagine synthetase Antibody, TS11 Antibody |
| Host | ASNS antibody was produced in Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Immunogen Species: | ASNS antibody was raised against Human |
| Antigen Type: | Recombinant protein |
| Immunogen: | ASNS antibody was raised against recombinant protein fragment contain a sequence corresponding to a region within amino acids 1 and 267 (P08243) of Asparagine synthetase. Percent identity by BLAST analysis: Human (100%); Bovine (95%); Porcine, Rat (94%); Mouse (93%). |
| Specificity: | Human ASNS |
| Reactivity: | Human |
| Purification: | Immunoaffinity purified |
| Presentation: | PBS, pH 7, 1% BSA, 20% Glycerol, 0.01% Thimerosal |
| Recommended Storage: | Aliquot and freeze at -20° C. Avoid freeze-thaw cycles. |
| Usage Summary: | Suggested starting dilutions are as follows: Western blot: 1:500-1:3000, IHC-Paraffin: 7.5 μg/ml. Not yet tested in other applications. Optimal working dilutions should be determined experimentally by the end user. |
| Uses: | IHC - Paraffin (7.5 μg/ml), Western blot (1:500 - 1:3000) (Optimal dilution to be determined by the researcher) |
| Size: | 100 μΙ |
| Concentration: | 0.6 mg/ml |
| | |

Immunohistochemistry Image:



Human Placenta: Formalin-Fixed, Paraffin-Embedded (FFPE)

Western Blot Image:



Sample (30 ug of whole cell lysate) A: K562 7.5% SDS PAGE ASNS antibody diluted at 1:1000

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences