



RNF216 / TRIAD3 Rabbit anti-Mouse Polyclonal (N-Terminus) Antibody - LS-B1430 - LSBio	
<b>CatalogID:</b>	LS-B1430
<b>Validation:</b>	This antibody replaces catalog number LS-C449. It has been validated for use in the following assays: IHC.
<b>Target:</b>	ring finger protein 216 (RNF216)
<b>Synonyms:</b>	RNF216 Antibody, TRIAD3 Antibody, UBCE7IP1 Antibody, Ring finger protein 216 Antibody, U711 Antibody, ZIN Antibody
<b>Family / Subfamily:</b>	Zinc Finger
<b>Host</b>	RNF216 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	RNF216 / TRIAD3 antibody was raised against Mouse
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	RNF216 / TRIAD3 antibody was raised against synthetic peptide from mouse RNF216 / TRIAD3.
<b>Specificity:</b>	peptide corresponding to 15 amino acids near the amino-terminus of mouse TRIAD3A
<b>Epitope:</b>	N-Terminus
<b>Reactivity:</b>	Mouse, Human
<b>Purification:</b>	Ion exchange chromatography
<b>Presentation:</b>	PBS, 0.02% sodium azide.
<b>Recommended Storage:</b>	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.
<b>Usage Summary:</b>	Immunohistochemistry: LS-B1430 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1430 was determined to be 5 ug/ml.
<b>Uses:</b>	IHC - Paraffin (5 µg/ml), Western blot (0.5 - 2 µg/ml) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**




Anti-RNF216 / TRIAD3 antibody IHC of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1430 concentration 5 ug/ml.

**Immunofluorescence Image:**



Immunofluorescence of TRIAD3A in Mouse Heart cells with TRIAD3A antibody at 20 ug/ml.

**Western Blot Image:**



Western blot of TRIAD3A in mouse heart cell lysates with TRIAD3A antibody at (A) 0.5, (B) 1, and (C) 2 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences