


WDHD1 Mouse anti-Human Monoclonal (C-Terminus) (20G10) Antibody - LS-B1656 - LSBio


CatalogID:	LS-B1656
Validation:	This antibody replaces catalog number LS-C41086. It has been validated for use in the following assays: IHC.
Target:	WD repeat and HMG-box DNA binding protein 1 (WDHD1)
Synonyms:	WDHD1 Antibody, AND-1 Antibody, CTF4 Antibody, AND1 Antibody, CHTF4 Antibody
Host	WDHD1 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2a
Clone Name:	20G10
Immunogen Species:	WDHD1 antibody was raised against Human
Immunogen:	WDHD1 antibody was raised against recombinant human WDHD1.
Specificity:	Recombinant (partial), C-terminal
Epitope:	C-Terminus
Reactivity:	Human, Mouse
Purification:	Affinity purified
Presentation:	Phosphate-buffered solution, pH 7.2, 0.09% sodium azide. Sourced in TCS.
Recommended Storage:	+4°C or -20°C, Avoid repeated freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B1656 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1656 was determined to be 15 ug/ml.
Uses:	IHC - Paraffin (15 µg/ml), Western blot (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:




Anti-WDHD1 antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1656 concentration 15 ug/ml.

Immunohistochemistry Image:



Anti-WDHD1 antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1656 concentration 15 ug/ml.

Western Blot Image:



Jurkat nuclear cell extracts (lane 1) and NIH3T3 nuclear extracts were resolved by electrophoresis, transferred to nitrocellulose, and probed with anti-AND-1 antibody (clone 20G10). Proteins were visualized using a goat anti-mouse-IgG secondary conjugated to HRP and chemiluminescence detection.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences