


NCOA1 / SRC-1 Mouse anti-Human Monoclonal (aa477-947) (1135) Antibody - LS-B1702 - LSBio	
<b>CatalogID:</b>	LS-B1702
<b>Validation:</b>	This antibody replaces catalog number LS-C20056. It has been validated for use in the following assays: IHC.
<b>Target:</b>	nuclear receptor coactivator 1 (NCOA1)
<b>Synonyms:</b>	NCOA1 Antibody, BHLHe42 Antibody, BHLHe74 Antibody, F-SRC-1 Antibody, Hin-2 protein Antibody, Nuclear receptor coactivator 1 Antibody, PAX3/NCOA1 fusion protein Antibody, KAT13A Antibody, Protein Hin-2 Antibody, Steroid receptor coactivator 1 Antibody, Steroid receptor coactivator-1 Antibody, NCoA-1 Antibody, RIP160 Antibody, SRC-1 Antibody, SRC1 Antibody
<b>Host</b>	NCOA1 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	1135
<b>Immunogen Species:</b>	NCOA1 / SRC-1 antibody was raised against Human
<b>Specificity:</b>	Amino acids 477-947 of human SRC-1.
<b>Epitope:</b>	aa477-947
<b>Reactivity:</b>	Human, Monkey, Mouse, Rat
<b>Purification:</b>	Protein G purified
<b>Presentation:</b>	PBS, pH 7.4.
<b>Recommended Storage:</b>	+4°C or -20°C, Avoid repeated freezing and thawing.
<b>Usage Summary:</b>	Immunohistochemistry: LS-B1702 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1702 was determined to be 5 ug/ml.
<b>Uses:</b>	IHC - Paraffin (5 µg/ml), Western blot, Immunoprecipitation (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**



Anti-NCOA1 antibody IHC of human brain, cerebellum. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1702 concentration 5 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences