


RNF213 Goat anti-Human Polyclonal (Internal) Antibody - LS-B1757 - LSBio	
CatalogID:	LS-B1757
Validation:	This antibody replaces catalog number LS-C47583. It has been validated for use in the following assays: IHC.
Target:	ring finger protein 213 (RNF213)
Synonyms:	RNF213 Antibody, C17orf27 Antibody, KIAA1618 Antibody, KIAA1554 Antibody, Mysterin Antibody, MYSTR Antibody, ALO17 Antibody, Moyamoya disease 2 Antibody, MYMY2 Antibody, NET57 Antibody, Ring finger protein 213 Antibody
Host	RNF213 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	RNF213 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	RNF213 antibody was raised against synthetic peptide C-KNPQTQTEERFRPQ from an internal region of human RNF213 (NP_065965.3). Percent identity with other species by BLAST analysis: Human (100%), Gorilla (93%).
Specificity:	Human RNF213.
Epitope:	Internal
Reactivity:	Human
Predicted Reactivity:	Gorilla
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B1757 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1757 was determined to be 1.25 ug/ml.
Uses:	IHC - Paraffin (1.25 µg/ml), ELISA (1:128000) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-RNF213 antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1757 concentration 25 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences