



ZMYM2 / RAMP Rabbit anti-Human Polyclonal (aa1250-1300) Antibody - LS-B1775 - LSBio	
CatalogID:	LS-B1775
Validation:	This antibody replaces catalog number LS-C47280. It has been validated for use in the following assays: IHC.
Target:	zinc finger, MYM-type 2 (ZMYM2)
Synonyms:	ZMYM2 Antibody, FIM Antibody, SCLL Antibody, MYM Antibody, Zinc finger MYM-type protein 2 Antibody, Zinc finger protein 198 Antibody, ZNF198 Antibody, Zinc finger, MYM-type 2 Antibody, RAMP Antibody
Host	ZMYM2 antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	ZMYM2 / RAMP antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	ZMYM2 / RAMP antibody was raised against synthetic peptide from human ZMYM2.
Specificity:	A portion of amino acids 1250-1300 of human Zinc finger protein 198
Epitope:	aa1250-1300
Reactivity:	Human, Orangutan, Monkey, Mouse, Rat, Bovine, Dog, Chicken
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.2% gelatin, 0.05% sodium azide.
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B1775 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1775 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 µg/ml), Western blot (0.1 - 1 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-ZMYM2 antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1775 concentration 10 ug/ml.

Western Blot Image:



Western blot of Zinc finger protein 198 in mouse skeletal muscle lysate in the 1) absence and 2) presence of immunizing peptide using antibody at 0.1 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences