


STRAP / MAWD Rabbit anti-Human Polyclonal (aa230-280) Antibody - LS-B1803 - LSBio


CatalogID:	LS-B1803
Validation:	This antibody replaces catalog number LS-C47371. It has been validated for use in the following assays: IHC.
Target:	serine/threonine kinase receptor associated protein (STRAP)
Synonyms:	STRAP Antibody, MAP activator with WD repeats Antibody, MAWD Antibody, PT-WD Antibody, UNR-interacting protein Antibody, UNRIP Antibody, WD-40 repeat protein PT-WD Antibody
Host	STRAP antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	STRAP / MAWD antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	STRAP / MAWD antibody was raised against synthetic peptide from human STRAP.
Specificity:	A portion of amino acids 230-280 of human STRAP
Epitope:	aa230-280
Reactivity:	Human, Chimpanzee, Monkey, Mouse, Bovine, Dog, Horse, Chicken, Xenopus, Zebrafish
Purification:	Protein G purified
Presentation:	PBS, 0.2% gelatin, 0.05% sodium azide.
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B1803 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1803 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 µg/ml), Western blot (1 - 2 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-STRAP antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1803 concentration 10 ug/ml.

Western Blot Image:



Western blot of human STRAP in Raw cell lysate in the 1) absence and 2) presence of immunizing peptide using antibody at 1 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences