


BAFF Mouse anti-Human Monoclonal (T7-241) Antibody - LS-B1909 - LSBio

CatalogID:	LS-B1909
Validation:	This antibody replaces catalog number LS-C41149. It has been validated for use in the following assays: IHC.
Target:	tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B)
Synonyms:	TNFSF13B Antibody, B-cell-activating factor Antibody, BAFF Antibody, ApoL related ligand TALL-1 Antibody, B-lymphocyte stimulator Antibody, CD257 Antibody, BLYS Antibody, Delta4 BAFF Antibody, Delta BAFF Antibody, TALL-1 Antibody, TALL1 Antibody, ZTNF4 Antibody, THANK Antibody, B lymphocyte stimulator Antibody, CD257 antigen Antibody, TNFSF20 Antibody
Family / Subfamily:	TNF / not assigned-TNF
Host	TNFSF13B antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Clone Name:	T7-241
Immunogen Species:	BAFF antibody was raised against Human
Immunogen:	BAFF antibody was raised against recombinant human TNFSF13B / BAFF.
Specificity:	Recombinant human BAFF(TL7) protein.
Reactivity:	Human
Purification:	Affinity purified
Presentation:	Phosphate-buffered solution, pH 7.2, 0.09% sodium azide. Sourced in Ascites.
Recommended Storage:	Store at 4°C. Do not freeze. Store undiluted.
Usage Summary:	Immunohistochemistry: LS-B1909 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1909 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 µg/ml), Flow Cytometry, ELISA (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-BAFF antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1909 concentration 10 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences