


NRM Rabbit anti-Human Polyclonal (aa200-250) Antibody - LS-B1979 - LSBio	
CatalogID:	LS-B1979
Validation:	This antibody replaces catalog number LS-C769. It has been validated for use in the following assays: IHC.
Target:	nurim (nuclear envelope membrane protein) (NRM)
Synonyms:	NRM Antibody, NRM29 Antibody, Nurim Antibody, Nuclear rim protein Antibody
Host	NRM antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	NRM antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	NRM antibody was raised against synthetic peptide from human NRM / Nurim.
Specificity:	A portion of amino acids 200-250 of human Nurim was used as the immunogen.
Epitope:	aa200-250
Reactivity:	Human, Mouse
Purification:	Protein G purified
Presentation:	PBS, 0.2% gelatin, 0.05% sodium azide.
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B1979 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B1979 was determined to be 5 ug/ml.
Uses:	IHC - Paraffin (5 μ g/ml), Western blot (1 - 3 μ g/ml) (Optimal dilution to be determine by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-NRM / Nurim antibody IHC of human small intestine. Immunohistochemistry of formalinfixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B1979 concentration 5 ug/ml.

