



DCAF6 / NRIP Rabbit anti-Human Polyclonal (aa800-850) Antibody - LS-B2046 - LSBio	
CatalogID:	LS-B2046
Validation:	This antibody replaces catalog number LS-C19347. It has been validated for use in the following assays: IHC-P.
Target:	DDB1 and CUL4 associated factor 6 (DCAF6)
Synonyms:	DCAF6 Antibody, ARCAP Antibody, PC326 Antibody, IQ motif and WD repeats 1 Antibody, RP4-745114.1 Antibody, IQWD1 Antibody, MSTP055 Antibody, NRIP Antibody
Host	DCAF6 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	DCAF6 / NRIP antibody was raised against Human
Specificity:	A portion of amino acids 800-850 of human NRIP was used as the immunogen.
Epitope:	aa800-850
Reactivity:	Human, Mouse, Rat, Dog
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.2% gelatin, 0.05% sodium azide.
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B2046 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2046 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 µg/ml), Western blot (1 - 3 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-DCAF6 / NRIP antibody IHC of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B2046 concentration 10 ug/ml.

Western Blot Image:



Western blot of NRIP in human muscle tissue lysate in the 1) absence and 2) presence of immunizing peptide, 3) mouse muscle tissue lysate and 4) rat muscle tissue lysate using antibody at 1 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences