



TNFRSF13B / TACI Rabbit anti-Human Polyclonal (aa116-132) Antibody - LS-B2094 - LSBio	
<b>CatalogID:</b>	LS-B2094
<b>Validation:</b>	This antibody replaces catalog number LS-C813. It has been validated for use in the following assays: IHC-P.
<b>Target:</b>	tumor necrosis factor receptor superfamily, member 13B (TNFRSF13B)
<b>Synonyms:</b>	TNFRSF13B Antibody, CD267 antigen Antibody, CD267 Antibody, CVID2 Antibody, TNFRSF14B Antibody, CVID Antibody, TACI Antibody
<b>Family / Subfamily:</b>	TNF Receptor / not assigned-TNF Receptor
<b>Host</b>	TNFRSF13B antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Immunogen Species:</b>	TNFRSF13B / TACI antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	TNFRSF13B / TACI antibody was raised against synthetic peptide from human TNFRSF13B / TACI.
<b>Specificity:</b>	A synthetic peptide corresponding to amino acids 116-132 of human TACI. Human and mouse TACI amino acid sequences are 65% conserved in the immunogen.
<b>Epitope:</b>	aa116-132
<b>Reactivity:</b>	Human, Mouse
<b>Purification:</b>	Protein G purified
<b>Presentation:</b>	PBS containing 0.05% sodium azide and 0.2% gelatin.
<b>Recommended Storage:</b>	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
<b>Usage Summary:</b>	Immunohistochemistry: LS-B2094 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2094 was determined to be 5 ug/ml.
<b>Uses:</b>	IHC - Paraffin (5 µg/ml), Western blot (5 µg/ml), Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	0.5 mg/ml

**Immunohistochemistry Image:**



Anti-TACI antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B2094 concentration 5 ug/ml.

**Western Blot Image:**



Western blot of 0.1 ug/ml of TACI in 10 ug/lane of Daudi lysate. A protein band of approximately 32 kDa is detected.

<b>Requested From:</b>	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 9/23/2014	
© 2014 LifeSpan BioSciences	