


BAFF Rat anti-Humar	Monoclonal (aa83-285) (Buffy-1) Antibody - LS-B2375 - LSBio
CatalogID:	LS-B2375
Validation:	This antibody replaces catalog number LS-C18640. It has been validated for use in the following assays: IHC-P.
Target:	tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B)
Synonyms:	TNFSF13B Antibody, B-cell-activating factor Antibody, BAFF Antibody, ApoL related ligand TALL-1 Antibody, B-lymphocyte stimulator Antibody, CD257 Antibody, BLYS Antibody, Delta4 BAFF Antibody, Delta BAFF Antibody, TALL-1 Antibody, TALL1 Antibody, ZTNF4 Antibody, THANK Antibody, B lymphocyte stimulator Antibody, CD257 antigen Antibody, TNFSF20 Antibody
Family / Subfamily:	TNF / not assigned-TNF
Host	TNFSF13B antibody was produced in Rat
Clonality:	Monoclonal
Isotype:	IgG2a
Clone Name:	Buffy-1
Immunogen Species:	BAFF antibody was raised against Human
Antigen Type:	Recombinant protein
Immunogen:	BAFF antibody was raised against recombinant human soluble BAFF (aa 83-285).
Specificity:	Recognizes membrane-bound human BAFF. Detects endogenous protein by FC.
Epitope:	aa83-285
Reactivity:	Human
Purification:	Gel filtration
Presentation:	PBS, 0.02% sodium azide.
Recommended Storage:	Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B2375 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2375 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (10 $\mu$ g/ml), ICC, Western blot, Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

## Immunohistochemistry Image:



Anti-BAFF antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody LS-B2375 concentration 10 ug/ml.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences