


CD163 Rabbit anti-Human Polyclonal (N-Terminus) Antibody - LS-B2661 - LSBio	
CatalogID:	LS-B2661
Validation:	This antibody replaces catalog number LS-C83693. It has been validated for use in the following assays: IHC-P.
Target:	CD163 molecule
Synonyms:	CD163 Antibody, CD163 antigen Antibody, Macrophage-associated antigen Antibody, M130 Antibody, CD163 molecule Antibody, Hemoglobin scavenger receptor Antibody, MM130 Antibody
Host	CD163 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	CD163 antibody was raised against Human
Immunogen:	CD163 antibody was raised against peptide derived from N-terminal sequence of human CD163.
Epitope:	N-Terminus
Reactivity:	Human
Purification:	Purified
Presentation:	20 mM Tris-HCl, pH 8.0, 20 mg/ml BSA, 0.05% sodium azide
Recommended Storage:	+4°C or -20°C, Avoid repeated freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B2661 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2661 was determined to be 20 ug/ml.
Uses:	IHC - Paraffin (20 μg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 μl
Concentration:	2 mg/ml

Immunohistochemistry Image:



Anti-CD163 antibody IHC of human liver. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody LS-B2661 concentration 20 ug/ml.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences