

PGR/PR/Progesterone Receptor Rabbit anti-Human Polyclonal (C-Terminus) Antibody - LS-B2983 - LSBio	
CatalogID:	LS-B2983
Validation:	This antibody replaces catalog number LS-C88867. It has been validated for use in the following assays: IHC-P.
Target:	progesterone receptor (PGR)
Synonyms:	PGR Antibody, NR3C3 Antibody, PR Antibody, Progesterone receptor Antibody
Family / Subfamily:	NHR / NR3 Steroid receptor
Host	PGR antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	PGR/PR/Progesterone Receptor antibody was raised against Human
Immunogen:	PGR/PR/Progesterone Receptor antibody was raised against peptide derived from C-terminus of inhibitory domain (IF) of human porgesterone receptor.
Specificity:	Synthetic peptide
Epitope:	C-Terminus
Reactivity:	Human
Purification:	Purified
Presentation:	20 mM Tris-HCl, pH 8.0, 20 mg/ml BSA, 0.05% sodium azide
Recommended Storage:	Store at -20°C. Aliquot to avoid freeze/thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B2983 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2983 was determined to be 20 ug/ml.
Uses:	IHC - Paraffin (20 μ g/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µl

Immunohistochemistry Image:

Anti-PGR / Progestero formalin-fixed, paraffin B2983 concentration 2	Receptor antibody IHC of human uterus. Immunohistochemistry of again.
Requested From:	Japan
	atory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
	Created on 9/23/2014
	© 2014 LifeSpan BioSciences