


MKI67 / Ki67 Rabbit anti-Human Polyclonal (C-Terminus) Antibody - LS-B2984 - LSBio	
CatalogID:	LS-B2984
Validation:	This antibody replaces catalog number LS-C88870. It has been validated for use in the following assays: IHC-P.
Target:	marker of proliferation Ki-67 (MKI67)
Synonyms:	MKI67 Antibody, Antigen KI-67 Antibody, Ki67 Antibody, KIA Antibody
Host	MKI67 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	MKI67 / Ki67 antibody was raised against Human
Immunogen:	MKI67 / Ki67 antibody was raised against peptide derived from C-terminal region of human Ki-67.
Specificity:	Synthetic peptide
Epitope:	C-Terminus
Reactivity:	Human
Purification:	Purified
Presentation:	20 mM Tris-HCl, pH 8.0, 20 mg/ml BSA, 0.05% sodium azide
Recommended Storage:	Store at -20°C. Aliquot to avoid freeze/thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B2984 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B2984 was determined to be 40 ug/ml.
Uses:	IHC - Paraffin (40 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	250 µl
Concentration:	1.5 mg/ml

Immunohistochemistry Image:



Anti-MKI67 / Ki-67 antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B2984 concentration 40 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences