



BAK1 / BAK Rabbit anti-Human Polyclonal (N-Terminus) Antibody - LS-B3333 - LSBio	
CatalogID:	LS-B3333
Validation:	This antibody replaces catalog number LS-C50599. It has been validated for use in the following assays: IHC-P.
Target:	BCL2-antagonist/killer 1 (BAK1)
Synonyms:	BAK1 Antibody, Apoptosis regulator BAK Antibody, BAK Antibody, BCL2L7 Antibody, Bcl-2-like protein 7 Antibody, BCL2-antagonist/killer 1 Antibody, Bcl2-L-7 Antibody, BCL2-like 7 protein Antibody, CDN1 Antibody, Pro-apoptotic protein BAK Antibody, BAK-LIKE Antibody
Host	BAK1 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	BAK1 / BAK antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	BAK1 / BAK antibody was raised against a synthetic peptide derived from N-terminal of human Bak.
Epitope:	N-Terminus
Reactivity:	Human, Mouse, Hamster
Purification:	Immunoaffinity purified
Presentation:	10mM PBS, pH 7.4, 1% BSA, 0.1% sodium azide
Recommended Storage:	Store at 2°C to 8°C degrees. Do not freeze.
Usage Summary:	Immunohistochemistry: LS-B3333 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B3333 was determined to be 10 ug/ml.
Uses:	IHC - Paraffin (1:300) (Optimal dilution to be determined by the researcher)
Size:	250 μΙ

## Immunohistochemistry Image:



Anti-BAK1 / BAK antibody IHC of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3333 concentration 10 ug/ml.

## Immunohistochemistry Image:



Anti-BAK1 / BAK antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3333 concentration 10 ug/ml.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences