


DAXX Rabbit anti-Human Polyclonal (aa261-274) Antibody - LS-B363 - LSBio	
CatalogID:	LS-B363
Validation:	This antibody replaces catalog number LS-C18933. It has been validated for use in the following assays: IHC.
Target:	death-domain associated protein (DAXX)
Synonyms:	DAXX Antibody, BING2 Antibody, CENP-C binding protein Antibody, DAP6 Antibody, Death-associated protein 6 Antibody, EAP1 Antibody, Fas-binding protein Antibody, HDaxx Antibody, ETS1-associated protein 1 Antibody
Family / Subfamily:	Apoptosis
Host	DAXX antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	DAXX antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	DAXX antibody was raised against synthetic peptide from human Daxx.
Specificity:	Amino acids 261-274 of human DAXX protein.
Epitope:	aa261-274
Reactivity:	Human, Mouse, Rat
Purification:	Immunoaffinity purified
Presentation:	0.02 M potassium phosphate, 0.15 M sodium chloride, pH 7.2, 0.01% sodium azide.
Recommended Storage:	+4°C or -20°C, Avoid repeated freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B363 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B363 was determined to be 2.5 ug/ml.
Uses:	IHC - Paraffin (2.5 µg/ml), Immunofluorescence, Western blot (1:200 - 1:2000), ELISA (1:7000 - 1:45000) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-DAXX antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B363 concentration 5 ug/ml.

Western Blot Image:



Anti-DAXX Antibody - Western Blot. Western blot of affinity purified anti-DAXX antibody shows detection of DAXX in RWPE-1 cell extracts. The membrane was probed with the primary antibody diluted to 1:200, and further probed with peroxidase conjugated anti-Rabbit IgG at a 1:25000 dilution. Personal Communication, Jie LIU, CCR-NCI, Bethesda, MD.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences